



národní
úložiště
šedé
literatury

Molecular description of *Blumeria graminis* f. sp. hordei isolates

Komínková, Eva; Malečková, Eva; Vanžurová, Hana; Dreiseitl, Antonín; Doležel, Jaroslav;
Valárik, Miroslav
2015

Dostupný z <http://www.nusl.cz/ntk/nusl-188958>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Licence Creative Commons Uveďte autora-Neužívejte dílo komerčně-Nezasahujte do díla 3.0 Česko

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 16.07.2018

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní nusl.cz .

Molecular description of *Blumeria graminis* f. sp. *hordei* isolates

Eva Komínková¹, Eva Malečková¹, Hana Vanžurová¹, Antonín Dreiseitl², Jaroslav Doležel¹
& Miroslav Valárik¹

¹Institute of Experimental Botany, Centre of the Region Haná for Biotechnological and Agricultural Research,
CZ-783 71 Olomouc, Czech Republic

²Agrotest Fyto Ltd., Havlíčkova 2787, CZ-767 01 Kroměříž, Czech Republic

The air-born fungus *Blumeria graminis* f. sp. *hordei* (*Bgh*) is a causal agent of barley powdery mildew. The pathogen attracts substantial attention due to its destructiveness. However, molecular diversity studies based on “house-keeping” genes do not provide sufficient resolution when applied to isolates from geographically limited regions. This study focused on developing a more efficient genotyping system capable to discriminate between closely related isolates. Whole genome sequence data were employed to design a panel of molecular markers based on microsatellites and insertion sites of transposable elements which represent an abundant part of the genome. A genotyping marker panel comprising 16 SSR, 14 SNP and 2 ISBP/RJM markers was applied on a set of 97 isolates originating from the Czech Republic, 50 Australian isolates and a collection of 11 isolates representing global *Bgh* diversity. The marker panel provided significant resolution of studied isolates, most of them showing unique genotype profiles. The analysis of phylogenetic relationship performed by neighbor-joining algorithm for 97 Czech isolates resulted in 87 separate clades and revealed high diversity of the pathogen population within a small geographical area. After supplementing with data on virulence of individual isolates, this study might open new opportunities of studying the host-pathogen relationship and patterns of the pathogen spatial distribution. This work has been supported by the Czech Ministry of Education, Youth and Sports (grant awards LD14105, LO1204).

