



národní  
úložiště  
šedé  
literatury

## **Inhibition of Benzothiophene HDS by Quinoline on PdMo and CoMo/Al<sub>2</sub>O<sub>3</sub> Catalysts**

Vít, Zdeněk  
2012

Dostupný z <http://www.nusl.cz/ntk/nusl-127030>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 30.09.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní [nusl.cz](http://nusl.cz) .

## INHIBITION OF BENZOTHIOPHENE HDS BY QUINOLINE ON PdMo AND CoMo/AL<sub>2</sub>O<sub>3</sub> CATALYSTS.

**Zdeněk Vít, Luděk Kaluža and Daniela Gulková**

*Institute of Chemical Process Fundamentals of the AS CR, v. v. i. ,  
Rozvojová 135, 165 02 Prague 6, Czech Republic*

Hydrodesulfurization (HDS) carried out industrially by means of conventional alumina supported CoMo catalysts is severely inhibited by basic nitrogen compounds present in the feed [1]. This is because these bases adsorb on the alumina support and also compete with sulfur reactants when interacting with active sites of the metal sulfide phase. This undesirable effect diminishes the performance of catalysts and becomes more important in hydrotreatment of heavier feeds with higher nitrogen content.

Recently, we observed that Mo/Al<sub>2</sub>O<sub>3</sub> sulfide catalysts promoted by Rh, Ru and Pd showed in simultaneous HDS of thiophene and HDN of pyridine higher HDS activities than conventional CoMo/Al<sub>2</sub>O<sub>3</sub> catalysts [2]. This effect was explained later by higher C-N bond cleavage activity of Rh and Ru promoted Mo catalysts which allows an easier removal of the adsorbed hydrogenated reaction intermediates [3].

In this work, the effect of quinoline on benzothiophene HDS was compared with the effect of pyridine on thiophene HDS. This is because benzothiophene and quinoline are closer to practical application. The main goal is to find, if the higher nitrogen tolerance observed during thiophene reaction occurs also with heavier sulfur and nitrogen molecules. Mo/Al<sub>2</sub>O<sub>3</sub> modified by Pd was selected for this study.

The support was  $\gamma$ -alumina (235 m<sup>2</sup>/g, Süd-Chemie AG). The catalysts contained 9 wt. % Mo and 2.0 % Co or 0.76 % Pd. The PdMo catalyst was prepared by impregnation of sulfided Mo/Al<sub>2</sub>O<sub>3</sub> by solution of Pd acetylacetonate. Catalysts were sulfided by 10 % H<sub>2</sub>S/H<sub>2</sub> at 400°C/1h. The benzothiophene HDS was carried out at 330°C/16 bar in the gas phase in an integral flow reactor with fixed bed of catalyst. The feed contained 1.8 wt.% S and 0, 500 or 1500 ppm of N. Decane was the solvent. Preliminary results up to now obtained are compared and discussed with relation to thiophene reaction.

### Acknowledgement

The support of the Czech Science Foundation (grant 104/09/0751) is gratefully acknowledged

### References:

- <sup>1</sup> Topsøe, H., Clausen, B.S., Massoth, F.E. in: J.R. Anderson and M. Boudart (Eds.), *Hydrotreating Catalysis, Science and Technology*, Springer, 1996, Vol. 11.
- <sup>2</sup> Vít, Z., Cinibulk, J., Gulková, D. *Appl. Catal. A* **2004**, 272, 99-107.
- <sup>3</sup> Vít, Z., Kaluža L. and Gulková D. *Top. Catal.* **2011**, 54, 1325-1330.

FTIR

J. I

New z  
connect  
catalysi  
to the  
The ot  
methyl  
by FTI  
isomorj  
alkylati

The Fl  
instrum  
by outj  
involve  
outgass  
The dis  
with a  
The alk  
under a  
using a

Brønste  
SSZ-33  
study d  
found  
Toluene  
[Fe]-SS  
to *p*-xy  
exhibits  
convers  
[Fe]-SS  
achieve  
iso/n-pr

Ackno

The aut

Refere

<sup>1</sup>Jones,  
471-48