



národní
úložiště
šedé
literatury

Application of HCSE Pulse Sequence to Detection of Small Signed Carbon-Carbon Coupling Constants

Blechta, Vratislav
2012

Dostupný z <http://www.nusl.cz/ntk/nusl-155593>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 06.05.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní nusl.cz .

APPLICATION OF HCSE PULSE SEQUENCE TO DETECTION OF SMALL SIGNED CARBON-CARBON COUPLING CONSTANTS.

Vratislav Blechta, Jan Schraml and Jan Sýkora

*Institute of Chemical Process Fundamentals of the ASCR, v. v. i.
Rozvojova 135, Prague 6 – Suchbát, 165 02, Czech Republic*

The HCSE pulse sequence [1] presented at previous Valtice conference aimed to detection of small ($J > 0.5$ Hz) homonuclear couplings. Originally this sequence was designed for silicon-silicon couplings.

The ability of HCSE to detect signed $^nJ(\text{C},\text{C})$ couplings, $n > 1$, is demonstrated on a set of 9 mono-substituted benzenes $\text{X}-\text{C}_6\text{H}_5$ with $\text{X} = \text{F}, \text{NO}_2, \text{MeO}, \text{Cl}, \text{Br}, \text{NO}, \text{N}\equiv\text{C}, \text{Me}$ and Me_3Si . All 36 $^2J(\text{C},\text{C})$ and $^3J(\text{C},\text{C})$ couplings in these compounds were calculated and compared with 26 experimental values in the paper by Witanowski et al. [2]. With the exception of one signed value all the reported [2] experimental values were absolute values only. The HCSE pulse sequence applied on these compounds yielded 34 signed values, 2 constants were not observed being probably very close to zero and therefore undetectable. The absolute values agree with the experimental ones given in ref. [2]. and all the detected signs agree with the computed ones. Influence of relaxations and coupling magnitudes on the signal intensities are discussed.

Financial support provided by The TACR (grant no. TA01010646) are gratefully acknowledged.

References

- [1] V. Blechta and J. Schraml, *Magn. Reson. Chem.*, **2012**, accepted DOI 10.1002/mrc.2869
- [2] M. Witanowski, K. Kamińska-Trela, Z. Biedrzycka, *J. Mol. Struct.*, **2007**, 844.