



národní  
úložiště  
šedé  
literatury

## **Effect of Adsorbable Foreign Gas on Evaporation of water from Porous Aerosol Particles**

Levdansky, V.V.  
2013

Dostupný z <http://www.nusl.cz/ntk/nusl-161337>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 28.09.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní [nusl.cz](http://nusl.cz) .

# EFFECT OF ADSORBABLE FOREIGN GAS ON EVAPORATION OF WATER FROM POROUS AEROSOL PARTICLES

Valeri LEVDANSKY<sup>1,2</sup>, Jiří SMOLÍK<sup>2</sup>, Vladimír ŽDÍMAL<sup>2</sup>

<sup>1</sup>Heat and Mass Transfer Institute NASB, Minsk, Belarus

<sup>2</sup>Institute of Chemical Process Fundamentals AS CR, v.v.i., Prague, Czech Republic

Keywords: aerosol particles, nanopores, evaporation

## INTRODUCTION

Adsorption of molecules of a foreign gas on the surface of nanoscale objects can influence phase transitions in the heterogeneous systems with nanoobjects. Some problems related to adsorption of foreign molecules on the surface of nanoscale particles (clusters) were considered in (Luijten et al., 1997; Levdansky et al., 2006). Here we discuss the effect of the adsorbable foreign gas on water evaporation from porous aerosol particles with nanoscale pores.

## RESULTS AND DISCUSSION

It is known that in a number of cases micro- and nanoscale pores can be inside solid aerosol particles (Fukuta, 1966). The aerosol particle in this case can be considered as the solid skeleton with fine (nanoscale) pores that are often filled with water (e.g. due to capillary condensation of the water vapor). The removal of water from the porous aerosol particles can be realized by water evaporation from pores, e.g. with increasing the particle temperature under the effect of solar radiation. Let us consider the influence of the adsorbable foreign gas on water evaporation from the fine cylindrical capillary in the free-molecular flow regime. Adsorption of molecules of a foreign gas on the water meniscus in the capillary can affect the surface tension of water as well as can block phase transitions on the meniscus surface. We consider for simplicity evaporation from the capillary with the fully wettable wall. The resulting flux of vapor molecules  $J$  passing through the capillary in water evaporation from the concave hemispherical meniscus can be written as

$$J = \frac{\pi d^2}{4} \frac{P_e}{(2\pi m k T)^{1/2}} Q W. \quad (1)$$

Here  $d$  is the diameter of the capillary that is assumed to be equal to the diameter of the meniscus,  $P_e$  is the saturated vapor pressure of water over the pure meniscus (without adsorbed foreign molecules),  $m$  is the mass of the water molecule,  $k$  is the Boltzmann constant,  $T$  is the temperature,  $W$  is the transmission probability of vapor molecules passing through the capillary, the value of  $Q$  in view of (Luijten et al., 1997) is given by

$$Q = \frac{1}{1 + K_{am} P_a} \left[ \left( \frac{1}{1 + K_{am} P_a} \right)^{-G} - H \right], \quad (2)$$

where  $G = 4Vn_0/d$ ,  $H = P_1/P_e$ ,  $n_0$  is the number density of molecules of the adsorbable foreign gas in the fully filled monolayer,  $P_1$  is the partial pressure of the water vapor at the exit end of the capillary,  $K_{am}$  is the Langmuir adsorption constant in adsorption of foreign molecules on the

meniscus,  $P_a$  is the partial pressure of the adsorbable foreign gas,  $V$  is the volume per molecule in the water.

Figure 1 shows the dependence of  $Q$  on  $\varphi_m = K_{am}P_a$ . It is seen from Fig. 1 that depending on  $H$  the value of  $Q$  in some range of  $\varphi_m$  can both decrease and increase with a rise in  $\varphi_m$ . Moreover, the introduction of the adsorbable foreign gas into the equilibrium at  $P_a = 0$  system can induce the resulting flux of water molecules (the line 3 in Fig. 1). It is also shown that adsorption of foreign molecules on the meniscus surface can affect the critical (equilibrium) diameter of the meniscus at which the resulting flux of water molecules on the meniscus surface is equal to zero.




Fig. 1.: Dependence of  $Q$  on  $\varphi_m$  in water evaporation from the cylindrical capillary at  $d = 10^{-8}$  m,  $T = 293$  K,  $n_0 = 6 \cdot 10^{18}$  m $^{-2}$  and different values of  $H$ ; 1:  $H = 0.93$ , 2:  $H = 0.97$ , 3:  $H = 1$ .

It is worth noting that the value of  $W$  in the general case can also depend on the pressure of the adsorbable foreign gas due to the effect of adsorbed foreign molecules on scattering of vapor molecules by the channel wall.

## CONCLUSIONS

It is shown that the influence of the adsorbable foreign gas on water evaporation from the porous aerosol particle with fine pores (capillaries) is related to a change in the flux of water molecules evaporating from the meniscus in the capillary as well as to a change in the transmission probability of water vapor molecules passing through the capillary due to the influence of adsorbed foreign molecules on scattering of water vapor molecules by the capillary wall.

## ACKNOWLEDGEMENTS

This work was supported by GAAVCR project IAA200760905.

## REFERENCES

- Fukuta N., Activation of atmospheric particles as ice nuclei in cold and dry air, *J. Atmos. Sci.*, 23, 741–750 (1966).
- Levdansky V.V., Smolík J., Moravec P., Influence of size effect and foreign gases on formation of nanoparticles, *Int. Commun. Heat Mass Transfer*, 33, 56–60 (2006).
- Luijten C.C.M., Bosschaart K.J., van Dongen M.E.H., High pressure nucleation in water/nitrogen systems, *J. Chem. Phys.*, 106, 8116–8123 (1997).