



národní  
úložiště  
šedé  
literatury

## **Inviscid Oscillations of Constrained Bubbles in Drops**

Vejražka, Jiří  
2013

Dostupný z <http://www.nusl.cz/ntk/nusl-170337>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 01.10.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní [nusl.cz](http://nusl.cz) .

# Inviscid oscillations of constrained bubbles and drops

Jiri Vejrázka, Lucie Vobecká, Jaroslav Tihon

Institute of Chemical Process Fundamentals, Prague, Czech Republic, vejrazka@icpf.cas.cz

In the instruments for measurements of the interfacial elasticity and viscosity, shape oscillations of the bubble or drop may occur when the drop is forced at higher frequencies. For better understanding of these shape oscillations, a linear inviscid theory is developed using variational principles<sup>1</sup> (via Lagrange's equations and Lagrange's lambda-multipliers). The theory allows determining eigenmodes (i.e. eigenfrequencies, eigenmode shapes and damping of eigenmode oscillations), but also response of the bubble shape to a motion of its support or to volume variations. Present theory covers also the cases previously analyzed by Strani and Sabetta<sup>2</sup> and Bostwick and Steen<sup>3</sup>. It can be applied to both bubbles and drops. Overall, the theory is flexible, as it can easily adapt to any type of constraints, and compared to previous treatments, it is also easy and intuitive.

The theory prediction has been compared to experiments. Good agreement is found for the case of small bubbles, which have spherical static shape. Experimental results for larger bubbles and drops deviate from the theory, as a neck is formed. It is shown that this deviation correlates well with a ratio of bubble volume to the volume at its detachment.




Figure: High-speed record of a bubble oscillating at a tip of capillary, a)  $D = 1.48$  mm, bubble oscillates mostly at first eigenmode, b)  $D = 1.19$  mm, oscillates mostly at third eigenmode

**Acknowledgement:** This research is supported by Ministry of Education, Youth and Sports (project no. LD13018).

<sup>1</sup> Vejrázka, J., Vobecká, L., Tihon, J. (2013): Linear oscillations of a supported bubble or drop. *Phys Fluids* **25**, 062102.

<sup>2</sup> Strani, M., Sabetta, F. (1984): Free vibrations of a drop in a partial contact with a solid support. *J Fluid Mech* **141**, 233.

<sup>3</sup> Bostwick, J. B., Steen, P. H. (1995): Capillary oscillations of a constrained liquid drop. *Phys Fluids* **21**, 032108.