



národní  
úložiště  
šedé  
literatury

## **A Class of Explicitly Solvable Absolute Value Equations**

Rohn, Jiří  
2014

Dostupný z <http://www.nusl.cz/ntk/nusl-170439>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 04.05.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní [nusl.cz](http://nusl.cz).



**Institute of Computer Science**  
**Academy of Sciences of the Czech Republic**

## **A Class of Explicitly Solvable Absolute Value Equations**

Jiří Rohn

<http://uivtx.cs.cas.cz/~rohn>

Technical report No. V-1202

04.02.2014



**Institute of Computer Science**  
**Academy of Sciences of the Czech Republic**

## **A Class of Explicitly Solvable Absolute Value Equations**

Jiří Rohn<sup>1</sup>


<http://uivtx.cs.cas.cz/~rohn>

Technical report No. V-1202

04.02.2014

Abstract:

We describe explicit formulae for the solution of a special case of absolute value equations.<sup>2</sup>



Keywords:

Absolute value equation, solution, explicit form.

---

<sup>1</sup>This work was supported with institutional support RVO:67985807.

<sup>2</sup>Above: logo of interval computations and related areas (depiction of the solution set of the system  $[2, 4]x_1 + [-2, 1]x_2 = [-2, 2]$ ,  $[-1, 2]x_1 + [2, 4]x_2 = [-2, 2]$  (Barth and Nuding [1])).

# 1 Introduction

Since its introduction in [3] ten years ago, the absolute value equation

$$Ax + B|x| = b$$

( $A, B \in \mathbb{R}^{n \times n}$ ,  $b \in \mathbb{R}^n$ ) has received quite a bit of attention. Various authors have proposed numerous methods for finding its solution (as e.g. via linear programming, nonlinear programming, etc.), but it seems that nobody has attempted so far to find an explicit form of the solution in some special cases. In this report we bring some result in this direction (Theorem 1). The assumption imposed on the right-hand side  $b$  may seem to be too restrictive, but even this partial result makes it possible to carry out the proof of the Hansen-Bliek-Rohn optimality result [2] in a new, more convincing fashion, which is possibly going to appear elsewhere.

Notation used:  $I$  is the identity matrix,  $e_k$  stands for the  $k$ th column of  $I$ , and the maximum of two vectors is taken entrywise.

## 2 The result

Our main result is formulated as follows.

**Theorem 1.** *Let  $\Delta \geq 0$ , let the matrix  $M = (I - \Delta)^{-1}$  satisfy*

$$M \geq I, \tag{2.1}$$

*and let there exist a  $k$  such that  $b_i \geq 0$  for each  $i \neq k$ . Then the equation*

$$x - \Delta|x| = b \tag{2.2}$$

*has a unique solution*

$$x = \max\{Mb, Mb - \frac{2(Mb)_k}{2M_{kk}-1}(M - I)e_k\}. \tag{2.3}$$

Moreover,

$$|x| = \max\{Mb, Mb - \frac{2(Mb)_k}{2M_{kk}-1}Me_k\}. \tag{2.4}$$

*Proof.* In view of nonnegativity of  $\Delta$ , the assumption (2.1) implies that  $\varrho(\Delta) < 1$ , hence  $\Delta^j \rightarrow 0$  as  $j \rightarrow \infty$ , so that the sequence  $\{x_i\}_{i=0}^\infty$  defined by  $x_0 = 0$  and

$$x_{i+1} = \Delta|x_i| + b \quad (i = 0, 1, \dots) \tag{2.5}$$

is Cauchian, thus convergent,  $x_i \rightarrow x$ . Taking the limit for  $i \rightarrow \infty$  in (2.5), we obtain

$$x = \Delta|x| + b, \tag{2.6}$$

hence  $x$  is a solution of (2.2). Assume that  $x'$  also solves (2.2). Then from

$$x - x' = \Delta(|x| - |x'|)$$

we obtain

$$|x - x'| \leq \Delta |x - x'|$$

and

$$(I - \Delta)|x - x'| \leq 0,$$

and premultiplying this inequality by the nonnegative matrix  $M$  leads to conclusion that

$$|x - x'| \leq 0,$$

hence  $x = x'$  and the solution  $x$  of (2.2) is unique.

Now, the assumption of nonnegativity of  $b_i$  for each  $i \neq k$  implies in the light of (2.6) that  $x_i \geq 0$  for  $i \neq k$ , so that  $|x_i| = x_i$  for each  $i \neq k$  and consequently we can write

$$|x| = x + (|x_k| - x_k)e_k. \quad (2.7)$$

Substituting this expression into (2.6), we obtain

$$(I - \Delta)x = (|x_k| - x_k)\Delta e_k + b$$

and by premultiplying by  $M$  we get

$$x = (|x_k| - x_k)(M - I)e_k + Mb \quad (2.8)$$

since  $M = (I - \Delta)^{-1}$  implies  $M\Delta = M - I$ . From this we can see that if  $x_k \geq 0$ , then

$$x = Mb.$$

If  $x_k < 0$ , then  $|x_k| = -x_k$  and from the  $k$ th equation in (2.8)

$$x_k = (|x_k| - x_k)(M_{kk} - 1) + (Mb)_k \quad (2.9)$$

we have

$$x_k = \frac{(Mb)_k}{2M_{kk} - 1} \quad (2.10)$$

(since  $M_{kk} \geq 1$  implies  $2M_{kk} - 1 \geq 1$ ), and finally from (2.8)

$$x = Mb - \frac{2(Mb)_k}{2M_{kk} - 1}(M - I)e_k.$$

Notice now that  $2M_{kk} - 1$  is positive and  $(M - I)e_k$  is nonnegative. Thus, if  $x_k \geq 0$ , then  $(Mb)_k \geq 0$  and

$$x = Mb \geq Mb - \frac{2(Mb)_k}{2M_{kk} - 1}(M - I)e_k,$$

hence

$$x = \max\{Mb, Mb - \frac{2(Mb)_k}{2M_{kk} - 1}(M - I)e_k\}, \quad (2.11)$$

and if  $x_k < 0$ , then  $(Mb)_k < 0$  (for otherwise (2.9) would imply  $x_k \geq 0$ ), hence

$$x = Mb - \frac{2(Mb)_k}{2M_{kk} - 1}(M - I)e_k \geq Mb,$$

so that (2.11) again holds. This proves (2.3).

To prove (2.4), we proceed in a similar way. If  $x_k \geq 0$ , then  $(Mb)_k \geq 0$  and

$$|x| = x = Mb \geq Mb - \frac{2(Mb)_k}{2M_{kk}-1}Me_k,$$

hence

$$|x| = \max\{Mb, Mb - \frac{2(Mb)_k}{2M_{kk}-1}Me_k\}, \quad (2.12)$$

whereas if  $x_k < 0$ , then  $(Mb)_k < 0$  and from (2.7), (2.10) we have

$$|x| = Mb - \frac{2(Mb)_k}{2M_{kk}-1}(M - I)e_k - \frac{2(Mb)_k}{2M_{kk}-1}e_k = Mb - \frac{2(Mb)_k}{2M_{kk}-1}Me_k \geq Mb,$$

which is again (2.12). This proves (2.4), and the proof is complete.  $\square$

We can also formulate the result in the following more explicit form.

**Theorem 2.** *Under assumptions and notation of Theorem 1, the unique solution  $x$  of the equation (2.2) is given by*

$$x = \begin{cases} Mb & \text{if } (Mb)_k \geq 0, \\ Mb - \frac{2(Mb)_k}{2M_{kk}-1}(M - I)e_k & \text{if } (Mb)_k < 0, \end{cases}$$

and its absolute value by

$$|x| = \begin{cases} Mb & \text{if } (Mb)_k \geq 0, \\ Mb - \frac{2(Mb)_k}{2M_{kk}-1}Me_k & \text{if } (Mb)_k < 0. \end{cases}$$

*Proof.* This is an immediate consequence of (2.3), (2.4).  $\square$

### 3 Application

With the help of Theorem 1 it is possible to derive closed-form formulae for matrices  $Q_z$  (introduced in [4]) for interval matrices of the form  $[I - \Delta, I + \Delta]$ , which then leads to a new proof of the Hansen-Blik-Rohn optimality result, as mentioned in the Introduction.

## Bibliography

- [1] W. Barth and E. Nuding, *Optimale Lösung von Intervallgleichungssystemen*, Computing, 12 (1974), pp. 117–125. [1](#)
- [2] J. Rohn, *Cheap and tight bounds: The recent result by E. Hansen can be made more efficient*, Interval Computations, 4 (1993), pp. 13–21. [1](#)
- [3] J. Rohn, *A theorem of the alternatives for the equation  $Ax + B|x| = b$* , Linear and Multilinear Algebra, 52 (2004), pp. 421–426. [1](#)
- [4] J. Rohn, *An algorithm for computing the hull of the solution set of interval linear equations*, Linear Algebra and Its Applications, 435 (2011), pp. 193–201. [3](#)