

Chiral Stationary Phases Based on Silica Modified by Helicenes

Bernard, Martin 2013

Dostupný z http://www.nusl.cz/ntk/nusl-170443

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 09.07.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní nusl.cz .

Chiral stationary phases based on silica modified by helicenes

Martin Bernard, Vladimír Církva, Jan Sýkora, Jan Storch*

Institute of Chemical Process Fundamentals of the ASCR, v. v. i., Rozvojova 2/135, CZ-165 02 Prague 6 – Suchdol, Czech Republic e-mail: bernard@icpf.cas.cz

Helicenes are inherently chiral o-condensed polyaromatic compounds with diverse potential applications such as chiral selectors in separation techniques. We have developed multigram-scale synthesis of various helicenes¹ based on photocyclization, some of them purposely derivatized² for anchoring on proper matrices via covalent bond. There are commercially available spherical stationary phases of defined mesh, possibly surface and pores are modified by appropriate reactive alkyl-group chains that commonly undergo further group modifications. Recently, we have run first trials potentially leading to stationary phases modified with racemic helicenes (preliminary results will be thoroughly discussed). After optimization of proper techniques and processes, the following step will be production of stationary phases modified with enantiomerically pure helicenes (scheme 1) and their testing for utilization in chiral HPLC.

Scheme 1

We appreciate the financial support from the Technology Agency of the Czech Republic (Grant No. TA01010646) and Ministry of Industry and Trade of the Czech Republic (Grant No. FR-TI3/628).

1. Storch J., Církva V., Bernard M., Vokál J., Pat. No. 303997/PV 2012 – 245, applied: 12.04.11.

2. Žádný J., Velíšek P., Jakubec M., Sýkora J., Církva V., Storch J., *Tetrahedron* **2013**, *69(30)*, 6213-6218.