



národní
úložiště
šedé
literatury

The Influence of Rutile Particles on Photo-induced Activity of the Sol-gel TiO₂/ITO Photoanode

Morozová, Magdalena
2014

Dostupný z <http://www.nusl.cz/ntk/nusl-175011>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 11.06.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní nusl.cz .

The influence of rutile particles on photo-induced activity of the sol-gel TiO₂/ITO photoanode

M. Morozova¹, P. Kluson¹, J. Krysa², O. Solcova¹

¹ Institute of Chemical Process Fundamentals of the ASCR, v. v. i., Rozvojová 135, 165 02 Prague 6, Czech Republic, tel. +420220390282, e-mail: morozova@icpf.cas.cz

² Institute of Chemical Technology, Technicka 5, 165 02 Prague 6, Czech Republic

Titanium dioxide (TiO₂) particles, mainly in anatase phase, immobilized to the thin layer form have found various applications in the gas sensors, photovoltaics, photocatalysis and photo-electrocatalysis due to its chemical and photo-chemical stability, as well as good photo-electrochemical properties. These applications require excellent charge separation and electron transport. It was found that the mixed crystallographic phase of anatase and rutile can exhibit a much higher photo-induced activity than the pure anatase or pure rutile form. The reason is that the photo-induced electrons can easily transfer from the anatase surface states to rutile, as well as from the anatase conduction band to rutile. These factors are responsible for better photocatalytic activity and for higher photo-current generated after illumination by the UV light. In this contribution the rutile effect on the structural and photo-electrochemical properties as well as on photocatalytical activity was studied.

Acknowledgement

The financial support of the Technology Grant Agency of the Czech Republic, projects No.: TA01020804 and TA03010548, is gratefully acknowledged.