



národní
úložiště
šedé
literatury

**Activity of Transition Metal Sulfides Supported on Alumina, Titania and Zirconia
in Parallel Hydrodesulfurization of 1-Benzothiophene and Hydrogenation of 1-Methyl-cyclohex**

Kaluža, Luděk
2014

Dostupný z <http://www.nusl.cz/ntk/nusl-177569>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 28.09.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní nusl.cz .

ACTIVITY OF TRANSITION METAL SULFIDES SUPPORTED
ON ALUMINA, TITANIA AND ZIRCONIA IN PARALLEL
HYDRODESULFURIZATION OF 1-BENZOTHIOPHENE
AND HYDROGENATION OF 1-METHYL-CYCLOHEX-1-ENE

Luděk Kaluža

*Institute of Chemical Process Fundamentals of the ASCR, v. v. i., Rozvojová 135,
165 02 Prague 6 – Suchbát, Czech Republic*

Sulfided conventional transition metals Co, Ni, Mo, and W and noble metals Rh, Pd, Ir, Pt, and Re deposited over conventional support gamma-Al₂O₃ (S_{HT} = 262 m²g⁻¹), and unconventional supports TiO₂ (anatase, S_{HT} = 140 m²g⁻¹) and ZrO₂ (baddeleyite, S_{HT} = 108 m²g⁻¹) were studied in parallel hydrodesulfurization of 1-benzothiophene (HDS) and hydrogenation of 1-methyl-cyclohex-1-ene (o-HYD) at 360 °C and 1.6 MPa. 1-Methyl-cyclohex-1-ene transformation was found to lead cyclic olefins containing also ethylcyclopentene and dimethylcyclopentene isomers. All volatile isomerization products were isolated directly from diluted reaction mixture via combination of chromatographic techniques including preparative gas chromatography, and were undoubtedly assigned by NMR spectroscopy [1]. Nevertheless, during each catalytic experiment, the model compound 1-methyl-1-cyclohex-1-ene yielded from more than 90% the mixture of the following four compounds: 1-, 3- and 4-methyl-cyclohex-1-enes and methylcyclohexane, which were quantified. It was ascertained that the 1-methyl-cyclohex-1-ene hydrogenation (o-HYD) that proceeded together with 1-benzothiophene hydrodesulfurization (HDS) did not significantly affected the HDS activity ranking of the studied transition metal sulfides supported on Al₂O₃, TiO₂ and ZrO₂ which was found before for the HDS reaction of 1-benzothiophene without 1-methyl-cyclohex-1-ene [2]. Mo, W, Co, and Ni sulfided catalysts exhibited relatively low activity in both HDS and o-HYD. For these sulfides, the TiO₂ and ZrO₂ support effect varied by the factor 0.4-9.3 and the o-HYD decreased the HDS activity by the factor 0.4-0.8. In contrast, the supported sulfides of Rh, Re, Pd, Pt, and Ir were highly active in HDS with low TiO₂ and ZrO₂ support effect (Rh/ZrO₂ being an exception) and o-HYD influenced the HDS activity by the factor 0.6-1.6. Nevertheless, Rh, Re, and Pd sulfided species exhibited relatively low activities in o-HYD reaction while Ir and Pt species exhibited high activities in o-HYD. The reference bimetallic CoMo and NiMo Al₂O₃ supported catalysts exhibited desirable high HDS but low o-HYD activities [3].

Acknowledgement. The author gratefully appreciates and acknowledges the financial support of the Czech Science Foundation (grant No. P106/11/0902).

References

- [1] Žáček P, Kaluža L, Karban J, Storch J, Šykora J (2014) React Kinet Mech Cat 112:335-346
 [2] Kaluža L, Zdrážil M (2008) Collect Czech Chem Commun 73:945-955
 [3] Kaluža L (2014) React Kinet Mech Cat (submitted).

AI DISTRIBUTION IN TNU-9 ZEOLITE

Robert Karz^{1,2}, Petr Klein¹, Veronika Pashkova¹, Štěpán Sklenák¹, Jiř

¹ J. Heyrovský Institute of Physical Chemistry, Academy of Sciences of the Republic, Dolejškova 2155/3, 182 23 Prague 8, Czech Republic

² J. Haber Institute of Catalysis and Surface Chemistry, Polish Academy of Sciences, Niezapominajek 8, 30-239 Cracow, Poland

Zeolites are crystalline microporous aluminosilicate molecular sieves defined, ordered framework formed by corner-sharing AlO₄ and SiO₄ tetrahedra of negatively charged AlO₄⁻ in the zeolite framework. This charge is balanced by redox catalytic reactions. The combination of a variety of tunable active sites and selectivity of the zeolite channel system and other zeolite properties result that zeolites represent the widest group of industrial heterogeneous catalysts.

Distribution of Al atoms in the zeolite framework determines its abundant active sites both for reactants and guest ions. This phenomenon was demonstrated for FER, MOR, BEA and ZSM-5 type zeolites [1-3]. Current work addresses the question of Al distribution within the framework of recently synthesized zeolite [4]. Methods used to solve presented problem encompass Vis-NIR, NMR spectroscopy as well as introduction of probe molecules into the zeolite framework. The composition of materials was analyzed using XRF. Samples of TNU prepared according to earlier described methods [4]. Post synthesis muller calcined in air and converted into sodium, ammonium or cobalt balanced forms.

Analysis of prepared samples resulted in ruling out the presence of Al sequences reflected in the ²⁹Si MAS NMR signal. Two main Al arrangements in the framework thus represent: (i) single Al atoms not able to stabilize Co²⁺ ions in the ring, able to accommodate bare Co²⁺ ions. Three types of Al pairs were identified in TNU-9 and their position in the channel system was discussed.

References:

- 1 Dědeček, J.; Wichterlová, B.; J. Phys. Chem. B **1999**, *103*, 1462-1476.
 2 Dědeček, J.; Kaucký, D.; Wichterlová, B.; Mic. Mes. Mat. **2000**, *35-36*, 483-487.
 3 Kaucký, D.; Dědeček, J.; Wichterlová, B.; Mic. Mes. Mat. **1999**, *31*, 73-87.
 4 Hong, S. B.; Min, H.-K.; Shin, C.-H.; Cox, P. A.; Warendorfer, S. J.; Weller, P. A. J. Am. Chem. Soc. **2007**, *129*, 10870-10885