



národní
úložiště
šedé
literatury

Preparation of Hydrodesulfurization Magnesia Supported Cobalt-Molybdenum Catalysts

Kaluža, Luděk
2014

Dostupný z <http://www.nusl.cz/ntk/nusl-177571>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 19.07.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní nusl.cz .

PREPARATION OF HYDRODESULFURIZATION MAGNESIA SUPPORTED
COBALT-MOLYBDENUM CATALYSTS

Luděk Kaluža, Daniela Gulková, Zdeněk Vít, Miroslav Zdražil

*Institute of Chemical Process Fundamentals of the ASCR, v. v. i., Rozvojová 135;
165 02 Prague 6 – Suchbát, Czech Republic*

Hydrodesulfurization (HDS) reactions are conventionally proceeded over gamma-Al₂O₃ supported CoMo, NiMo or NiW sulfides. Alternative supports are intensively studied to increase activity in HDS to meet increasingly stringent regulation on sulfur level in fuels. Particularly, MgO support exhibits higher basicity in comparison to gamma-Al₂O₃, which results in high dispersion and activity of deposited MoS₂, high synergy in CoMo phase and high resistance to coking. These advantages are still insufficiently utilized mainly because of the difficulties with the deposition of CoMo active phase onto hydrothermally little stable surface of MgO. However, we have reported on reaction of (NH₄)₂MoO₇ or MoO₃ with the high surface area MgO in methanol slurry which resulted in well-defined saturated Mo monolayer and high activity in HDS of 1-benzothiophene. In the second step, Co was deposited over MoO₃/MgO catalyst from the solution of Co(NO₃)₂ in methanol by impregnation, which resulted in high synergic effect of Co and about 2-fold increase in activity in benzothiophene HDS than it was observed for conventional gamma-Al₂O₃ support. The aim of the present work was to elucidate the possibility of deposition of molybdenyl- and cobalt-acetylacetonates on the high surface area MgO by impregnation from solutions of these compounds in methanol. Furthermore, non-aqueous solutions of MoO₃ and CoCO₃, (NH₄)₂MoO₇ and Co(NO₃)₂, MoO₃(C₂H₅O)₂ and Co(C₂H₅O)₂ with chelating agent, nitriofacetic acid (NTA) were studied for CoMo deposition in one impregnation step. It was found that the highly dispersed and x-ray amorphous loadings of Mo linearly corresponded to specific surface area of the MgOs studied (about 330 m² g⁻¹ and 500 m² g⁻¹) and HDS activities. TPR measurement revealed that high Mo content in the Mo/MgO catalyst led to a decrease in the temperature of sulfides reduction and decrease in hydrogen consumption. The decrease of the reduction temperature is well related to the increase in the catalytic activity. In contrast, the decrease in hydrogen consumption indicated a decrease of sulfur edge sites, which may be in agreement with the high loadings of Mo but it is contradicted to the high HDS activities observed. Neither the hydrogen consumption nor the O₂ uptake, as a measure of the sulfide dispersion, explained the HDS activity trends. Furthermore, the depositions of Co resulted in 13-46 fold promotion of the activity of Mo/MgO catalysts. This promotion, however, was highly sensitive to the method of Co deposition and the MgO type. The most active catalysts were prepared by i) deposition of Co onto oxidic Mo catalyst prepared from MoO₃/methanol slurry or ii) deposition of Co onto presulfided Mo catalyst prepared from MoO₃(C₂H₅O)₂/methanol solution and iii) by co-impregnation from MoO₃(C₂H₅O)₂, Co(C₂H₅O)₂ and NTA solution of freshly calcined and dechlorinated MgO support with surface area of about 330 m² g⁻¹. The authors gratefully acknowledge the financial support of the Czech Science Foundation (grant no. P106/11/0902).

Reference

L. Kaluža, D. Gulková, Z. Vít, M. Zdražil, Appl. Catal. B: Environ. 162 (2015) 430-436

CONDENSATION REACTION CATALYZED BY FUNCTIONALIZED
MCM-41

Eva Vrbková, Eliška Vyskočilová, Libor Červený

*Department of organic technology, Institute of chemical technology, Technická 5,
Praha 6, CZ 166 28, Czech Republic*

Condensation reactions play an important role in the synthesis of fine chemicals *f. ex.* containing double bond in conjugation with carbonyl group. These reactions are usually catalyzed by homogeneous basic catalysts or by homogeneous catalyst containing metals. With increasing interest in environment this catalysts are replaced by heterogeneous catalysts - offering easy separation from reaction mixture and reuse.

Within this work, several types of functionalized MCM-41 were prepared using post-grafting method - MCM-41 was grafted by 3-propylcarboxy-, 3-propylsulfoxy-, 3-aminopropyl-, 3-(2-aminoethyl)-3-aminopropyl-, 3-glycidoxypropyl and 3-propylisocyanate groups.

Prepared functionalized MCM-41s were used as catalysts in following reactions:

1. Aldol condensation of benzaldehyde and heptanal, which gives desired product *jasminaldehyde*.
2. Acylation of syringaldehyde using acetic anhydride - desired product *syringaldehyde diacetate*.
3. Acylation reaction of ethylvanillin using acetic anhydride - desired product *ethylvanillin diacetate*.
4. Acylation of 4-dimethylaminobenzaldehyde using acetic anhydride - desired product *4-dimethylaminobenzaldehyde diacetate*.

Influence of the reactants amount, temperature, type of catalyst or solvent on the reaction course was monitored. Optimal reaction conditions were suggested.

Acknowledgement

Financial support from specific university research (MSMT No 20/2014).