



národní  
úložiště  
šedé  
literatury

## **Seasonal Differences in Chemical Composition of Atmospheric Aerosol Studied with High Resolution in Prague**

Kubelová, Lucie  
2015

Dostupný z <http://www.nusl.cz/ntk/nusl-189279>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 05.05.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní [nusl.cz](http://nusl.cz) .

# Seasonal differences in chemical composition of atmospheric aerosol studied with high resolution in Prague

L. KUBELOVÁ<sup>12\*</sup>, P. VODIČKA<sup>1</sup>, O. MAKEŠ<sup>12</sup>, J. SCHWARZ<sup>1</sup>  
AND V. ŽDÍMAL<sup>1</sup>

<sup>1</sup>Institute of Chemical Process Fundamentals of the CAS, Prague, Czech Republic (\* kubelova@icpf.cas.cz)

<sup>2</sup>Institute of Environmental Studies, Faculty of Science, Charles University, Prague, Czech Republic

The compact-Time of Flight-Aerosol Mass Spectrometer (AMS) was used to describe seasonal differences in atmospheric aerosol at a Prague suburban site. The data were compared with measurements done with other instruments such as Sunset organic and elemental carbon analyzer, TSI Scanning Mobility Particle Sizer (SMPS), and results from ion chromatography, and with meteorological model.

In winter, we observed higher total concentration of NR-PM1 (non-refractory PM1) than in summer. Furthermore, the maxima of size distributions were higher in winter and during polluted episodes than in summer and during clean episodes. We discussed the influence of different seasonal sources, e.g. heating in winter. The daily cycles of the main AMS compounds were strongly influenced by the dilution effect of the mixing layer. The daily cycle of sulphate anion was influenced by the transport from the upper boundary layer, i.e. reservoir effect, and by the oxidation of SO<sub>2</sub> to SO<sub>4</sub><sup>2-</sup> (Fig.1).

A comparison of the organic fragments f43 and f44 [1] revealed less photochemically aged aerosol in winter compared to summer. Also, the winter aerosol was more influenced by biomass burning.




Fig.1: Daily cycle of (a) SO<sub>2</sub> and (b) sulfate anion in summer at a suburban measurement site Prague Suchdol.

[1] Ng *et al.* (2010) *Atmos. Chem. Phys.*, **10**, 4625–4641.

Ack: CSF P209/11/1342.