



národní
úložiště
šedé
literatury

Highly Time-Resolved Aerosol Measurement at a Suburban Site in Prague

Kubelová, Lucie
2015

Dostupný z <http://www.nusl.cz/ntk/nusl-200958>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 24.07.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní [nusl.cz](http://www.nusl.cz) .

HIGHLY TIME-RESOLVED AEROSOL MEASUREMENT AT A SUBURBAN SITE IN PRAGUE

Lucie KUBELOVÁ^{1,2}, Petr VODIČKA¹, Jaroslav SCHWARZ¹, Otakar MAKEŠ^{1,2}, Vladimír ŽDÍMAL¹

¹ Institute of Chemical Process Fundamentals of the CAS, v.v.i., Prague, Czech Republic, kubelova@asuch.cas.cz

² Department of Environmental Studies, Faculty of Science, Charles University, Prague, Czech Republic, kubelovl@natur.cuni.cz

Keywords: Atmospheric aerosols, Chemical composition, AMS, PM1

INTRODUCTION

Highly time-resolved measurements of atmospheric aerosols are important as they enable us to elucidate the aerosol sources and lifecycle processes (Zhang, 2011). In our study we carried out two six-week measurements of submicron fraction of non-refractory atmospheric aerosol. The measurement was performed with 1-minute resolution and analysed with respect to influence of meteorological conditions. The measurement was done as a part of a pan-European project ACTRIS.

EXPERIMENTAL SETUP

The measurements were done at Prague Suchdol suburban measurement site, which is located approximately six kilometers north west from the Prague city center. During the two measurement campaigns (summer 2012, winter 2013), we deployed the c-ToF-AMS, field Organic Carbon/Elemental Carbon OC/EC analyser and PM1 filter measurement analysed by Ion Chromatography (IC). The c-ToF-AMS provides us with time resolved chemical composition and size distribution of aerosol particles (Drewnick, 2005). The vaporization and ionization occurred at 600°C and 70eV, respectively.

Tab.1: Average composition of non-refractory submicron aerosol during summer and winter campaigns in Prague Suchdol.

Compound	SUMMER		WINTER	
	AVERAGE ± ST.DEV. ($\mu\text{g}/\text{m}^3$)	SHARE	AVERAGE ± ST.DEV. ($\mu\text{g}/\text{m}^3$)	SHARE
Org	4.2 ± 3.2	51.2%	8.4 ± 6.9	39.5%
NH ₄ ⁺	1.2 ± 0.9	14.0%	2.8 ± 2.1	13.1%
SO ₄ ²⁻	2.0 ± 1.6	24.4%	4.4 ± 3.7	20.9%
NO ₃ ⁻	0.8 ± 0.9	9.7%	5.4 ± 4.6	25.4%
Cl ⁻	0.1 ± 0.1	0.7%	0.23 ± 0.26	1.1%
Total	8.3 ± 6.0	x	21.2 ± 16.4	x

RESULTS AND CONCLUSIONS

Table 1 shows the average composition of the summer and winter campaign. In winter, there was a significantly higher share of nitrate than in summer, which we explain by the influence of lower temperatures promoting the particulate phase. From comparison with calculations done using the HYSPLIT model, we found an inverse relationship between the mixing height of the boundary layer and the overall level of pollution. There was also a strong correlation between the arrival of continental (maritime) airmasses and an increased (decreased) level of pollution. The analysis of organic fragments f43, f44, and f60 confirmed the influence of domestic heating on winter aerosol. The high time-resolution enabled us to study the daily cycles in detail. The daily cycles were influenced by the mixing height of the boundary layer, photochemical reactions and entrainment of pollution from higher atmospheric layers.

ACKNOWLEDGEMENT

This project has received funding from the EU Horizon 2020 Research and Innovation Programme under Grant Agreement No. 654109 and from the Czech Science Foundation under project No. CSF P209/11/1342.

REFERENCES

- Drewnick, F., Hings, S., DeCarlo, P.F., Jayne, J.T., Gonin, M., Fuhrer, K., Weimer, S., Jimenez, J.L., Demerjian, K.L., Borrmann, S., Worsnop, D.R., A new time-of-flight aerosol mass spectrometer (TOF-AMS) - Instrument description and first field deployment, *Aerosol Sci. Technol.*, 39, 637-658, (2005).
- Kubelová L., Vodička P., Schwarz J., Cusack M., Makeš O., Ondráček J., Ždímal V., A study of summer and winter highly time-resolved submicron aerosol composition measured at a suburban site in Prague, *Atmospheric Environment*, 118, 45-57, (2015).
- Zhang Q., Jimenez J. L., Canagaratna M. J., Ulbrich I. M., Nga L. N., Worsnop D. R., Sun Y., Understanding atmospheric organic aerosols via factor analysis of aerosol mass spectrometry: a review, *Aerosol Analysis*, 401, 3045-3067, (2011).