



národní
úložiště
šedé
literatury

Comparison of Atmospheric Aerosol Volatility at a Suburban and Rural Measurement Site.

Kubelová, Lucie
2016

Dostupný z <http://www.nusl.cz/ntk/nusl-261497>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 28.04.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní nusl.cz .

COMPARISON OF ATMOSPHERIC AEROSOL VOLATILITY AT A SUBURBAN AND RURAL MEASUREMENT SITE

Lucie KUBELOVÁ^{1,2}, Petr VODIČKA¹, Jaroslav SCHWARZ¹, Otakar MAKEŠ^{1,2},
Vladimír ŽDÍMAL¹

¹Institute of Chemical Process Fundamentals of the CAS, v.v.i., Prague, Czech Republic,
kubelova@asuch.cas.cz

²Department of Environmental Studies, Faculty of Science, Charles University, Prague,
Czech Republic, kubelovl@natur.cuni.cz

Keywords: Atmospheric aerosols, AMS, PM1, Volatility

INTRODUCTION

One of the most important characteristics of atmospheric aerosols is their volatility. Volatility affects the formation, life time, and removal of aerosol particles (Huffman et al., 2009). Accurate description of volatility of atmospheric aerosols can lead to a better identification of their sources and it is also crucial for models incorporating condensation and aging of aerosols (Wu et al., 2009; Zhao et al., 2016). However, there is still a lack of knowledge regarding aerosol properties such as volatility especially in case of organic aerosols (Lee et al., 2010).

To increase our understanding of aerosols volatility, we have performed summer (21.8.-2.10.2012; 17.6.-18.8.2014) and winter (19.2.-28.5.2013; 9.1.-13.3.2014) measurement campaigns with a thermodenuder at a suburban site in Prague Suchdol and at a rural site Košetice, respectively.

EXPERIMENTAL SETUP

The following instruments were deployed during the summer and winter measurement campaigns with a thermodenuder: Compact Time-Of-Flight Aerosol Mass Spectrometer (C-ToF-AMS), Scanning Mobility Particle Sizer (SMPS), Filter Measurement Analyzed by Ion Chromatography (IC).

The collection efficiencies for the non-thermodenuded sample measured by the AMS during particular campaigns were determined by comparison of sulphate concentrations measured by the AMS and chemical analyses of PM1 filter samples taken in parallel. The resulting values are following: Suchdol Summer (PS) 0.36, Suchdol Winter (PW) 0.33, Košetice Summer (KS) 0.25*, Košetice Winter (KW) 0.14*.

* Uncorrected for sample dilution.

RESULTS

Figure 1 shows the ratio of thermodenuded and non-thermodenuded samples for particular campaigns. Sulphate was the least volatile compound during all campaigns, followed by ammonium, chloride, organics and nitrate - the most volatile compounds. The relatively lower volatility of nitrate in summer compared to winter is caused by higher average temperatures in summer leading to its evaporation prior to entering the sampling head. We have also studied the volatility in relationship to the origin of relevant air masses.




Fig. 1: Ratio of thermodenuded and non-thermodenuded mass concentrations measured by the AMS during particular campaigns

ACKNOWLEDGEMENT

This project has received funding from the European Union Horizon 2020 Research and In-novation Programme under Grant Agreement No. 654109, from the Czech Science Foundation under project No. CSF P209/11/1342 and also from the Ministry of Education, Youth and Sports under project Nr. LM2015037.

REFERENCES

- Huffman, J.A., Docherty, K.S., Mohr, C., Cubison, M.J., Ulbrich, I.M., Ziemann, P.J., Onasch, T.B., Jimenez, J.L. Chemically-resolved volatility measurements of organic aerosol from different sources. *Environ. Sci. Technol.*, 43, 5351–5357, (2009).
- Lee, B.H., Kostenidou, E., Hildebrandt, L., Riipinen, I., Engelhart, G.J., Mohr, C., Decarlo, P.F., Mihalopoulos, N., Prevot, A.S.H., Baltensperger, U., Pandis, S.N. Measurement of the ambient organic aerosol volatility distribution: Application during the Finokalia Aerosol Measurement Experiment (FAME-2008). *Atmos. Chem. Phys.*, 10, 12149–12160, (2010).
- Wu, Z., Poulain, L., Wehner, B., Wiedensohler, A., Herrmann, H., Characterization of the volatile fraction of laboratory-generated aerosol particles by thermodenuder-aerosol mass spectrometer coupling experiments. *J. Aerosol Sci.*, 40, 603–612, (2009).
- Zhao, B., Wang, S., Donahue, N.M., Jathar, S.H., Huang, X., Wu, W., Hao, J., Robinson, A.L. Quantifying the effect of organic aerosol aging and intermediate-volatility emissions on regional-scale aerosol pollution in China. *Sci. Rep.*, 6, 28815, (2016).