

Revised Synthesis and Derivatisation of 2-Bromo[6]helicene.

Jakubec, Martin 2016

Dostupný z http://www.nusl.cz/ntk/nusl-261724

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 21.05.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní nusl.cz .

Revised Synthesis and Derivatisation of 2-Bromo[6]helicene

Martin Jakubec¹, Tomáš Beránek¹, Pavel Jakubík¹, Jan Sýkora¹, Jaroslav Žádný¹, Jan Storch¹

¹Institute of Chemical Process Fundamentals, v.v.i., Rozvojová 1/135, Prague 6, 165 02 e-mail: jakubecm@icpf.cas.cz

2-Bromo[6]helicene¹ represents a desirable starting material for preparation of a variety of different derivatives bearing helicene-2-yl moiety. We developed a revised shorter and more efficient synthesis.

We also present robust procedures for preparation of several derivatives, which include products of carbon – carbon coupling reactions, such as Suzuki-Miyaura reaction, Heck reaction or Rosemund-von Braun reaction. Other heteroatoms, such as boron, nitrogen, oxygen, sulphur and phosphorus were also introduced into the structure with moderate to good yields. Most of the reactions were performed in a microwave reactor, keeping the reaction time to a necessary minimum. In general, this methodology enables the introduction of various functional groups and allows for the utilization of helicenes in various fields of research, namely material chemistry, supramolecular chemistry, or optoelectronics².

For the financial support, the Czech Science Foundation (Project No. 15-12719S) is gratefully acknowledged.

1 Lightner, D. A.; et. al. J. Am. Chem. Soc. 1972, 94 (10), 3492.

2 Gingras, M. Chem. Soc. Rev. 2013, 42, 1051.