



národní  
úložiště  
šedé  
literatury

**Efficient Synthesis of Anion Receptors Based on the Phenoxythiin Macrocycle.**

Landovský, T.  
2016

Dostupný z <http://www.nusl.cz/ntk/nusl-261725>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 28.09.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní [nusl.cz](http://www.nusl.cz) .

## Efficient Synthesis of Anion Receptors Based on the Phenoxathiin Macrocycle

Tomáš Landovský,<sup>1</sup> Petra Cuřínová,<sup>2</sup> Václav Eigner,<sup>3</sup> Jan Budka,<sup>1</sup> Pavel Lhoták\*,<sup>1</sup>

<sup>1</sup>Department of Organic Chemistry, University of Chemistry and Technology Prague, Technická 5, Prague 6, Czech Republic, 166 28


<sup>2</sup>Institute of Chemical Process Fundamentals of the AS CR, v.v.i., Rozvojová 2/135, Prague 6, Czech Republic, 165 02

<sup>3</sup>Department of Solid State Chemistry, University of Chemistry and Technology Prague, Technická 5, Prague 6, Czech Republic, 166 28  
e-mail: landovst@vscht.cz

Thiacalixarenes are macrocyclic compounds that are widely used in supramolecular chemistry. The main advantage of their skeleton is the possibility of its simple modifications.<sup>1</sup> Being one of the easily accessible derivatives, phenoxathiin macrocycle is interesting due to its inherent chirality.<sup>2</sup> However, wider utilization of this novel compound is problematic because of the lack of effective derivatization techniques.<sup>3</sup>

We present the first and very efficient way how to introduce the amino groups to the upper rim of the macrocycle. It was performed by using the *ipso*-nitration of the fully S-oxidized macrocycle followed by selective reduction of nitro groups. The structures of most of the products were confirmed by X-ray crystallography of monocrystals.

Resulting amino derivative was further reacted with two selected isocyanates, providing the ureido-type receptors (Scheme 1). Their abilities to complex common anions were measured by NMR titration and unexpectedly high association constants were observed in all cases.



Scheme 1

This work was supported by Czech Science Foundation (reg. No. 16-13869S).

1 Kumar, R.; Lee, Y. O.; Bhalla, V.; Kumar, M.; Kim, J. S. *Chem. Soc. Rev.* **2014**, *43*, 4824.

2 Polívková, K.; Šimánová, M.; Budka, J.; Cuřínová, P.; Císařová, I.; Lhoták, P. *Tet. Lett.* **2009**, *50*, 6347.

3 Vrzal, L.; Kratochvílová-Šimánová, M.; Landovský, T.; Polívková, K.; Budka, J.; Dvořáková, H.; Lhoták, P. *Org. Biomol. Chem.* **2015**, *13*, 9610.