



národní
úložiště
šedé
literatury

Ruthenium (II) Complexes with 6,7-Diaza[5]phenacene.

Beránek, Tomáš
2016

Dostupný z <http://www.nusl.cz/ntk/nusl-261727>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 10.04.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní nusl.cz.

Ruthenium(II) Complexes with 6,7-Diaza[5]phenacene

Tomáš Beránek,¹ Jaroslav Žádný,^{*,1} Pavel Jakubík,¹ Vladimír Církva,¹ Jan Storch¹


¹*Institute of Chemical Process Fundamentals, v.v.i., Rozvojová 1/135,
Prague 6, 165 02
e-mail: beranekt@icpf.cas.cz*

Ruthenium complexes with polydentate-nitrogen ligands are widely used in the field of photocatalysis¹, optoelectronics² or as chromophors in dye-sensitized solar cells².

In this work we focused on preparation of novel bidentate and fully aromatic *N,N*-ligand and its ruthenium complexes. 6,7-Diaza[5]phenacene (**1**) was synthetized from 4-bromoisoquinoline by microwave-assisted tandem Hyama-Heck cross coupling reaction³ followed by photocyclization (Scheme 1).

With 6,7-diaza[5]phenacene (**1**) in our hand, we prepared some new ruthenium(II) complexes (e.g. **2**), characterized them from structural point of view and determined some of their electronic and optical properties with regard to their application potential.

Scheme 1



This work was supported by the Technology Agency of the Czech Republic (Grant No. TA01010646).

1 Prier, Ch.; Rankic, D.; MacMillan, D. *Chem. Rev.* **2013**, *113*, 5322.

2 Hiroki M., *Kobunshi* **2014**, *63*, 866.

3 Grätzel M., et al. *J. Am. Chem. Soc.* **1985**, *107*, 2988.

4 Gordillo A., et. al. *J. Am. Chem. Soc.*, **2013**, *135*, 13749.