



národní
úložiště
šedé
literatury

Thermodynamic Properties of Saccharinate-based Ionic Liquids: Comparison of Experimental Data to a Prediction by Means of COSMO-RS.

Čanji, Maja
2017

Dostupný z <http://www.nusl.cz/ntk/nusl-320489>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 05.05.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní nusl.cz .

Thermodynamic Properties of Saccharinate-based Ionic Liquids: Comparison of Experimental Data to a Prediction by means of COSMO-RS

Student: Maja Čanji, MSc.

Supervisor: Ing. Magdalena Bendova, Ph. D.

As an overall general aim of this study, thermodynamic properties of hydrophilic 1-alkyl-3-methylimidazolium-based ILs ($n = 4, 6, 8, 10, 16$) with a saccharinate (Sac^-) anion have been investigated. Saccharinate is a natural nontoxic anion, contributing to the application potential of the present ILs series. They were therefore used in a previous study for a laboratory-scale extraction of glaucine from its plant.¹ To design an industrial-scale extraction, it is important to have a good knowledge of the physico-chemical properties in the relevant mixtures of ILs with molecular solvents.

In the present contribution, density of 1-alkyl-3-methylimidazolium saccharinates ($n = 4, 6, 8, 10$) and isobaric heat capacity for 1-butyl-3-methylimidazolium saccharinate were measured in this work for a basic characterization of the studied ionic liquids. Modelling based on the COSMO-RS² model was used to predict density and heat capacity³ for the studied ILs and compared with experimental data. Moreover, a prediction by means of COSMO-RS of liquid-liquid equilibria⁴ for 1-decyl-3-methylimidazolium saccharinate with water was carried out and compared with a preliminary experiment of its limited miscibility.

References

1. Bogdanov. G.M.; Svinyarov. I.; Keremedchieva. R.; Sidjimov. A. Ionic liquid supported solid-liquid extraction of bioactive alkaloids I. New HPLC metod for quantitative determination of glaucine in *Glaucium flavum* Cr. (*Papaveraceae*). *Sep. Pur. Tech.* **2012**, *97*, 221–227.
2. Klamt, A. A Conductor-like screening model for real solvent: A new approach to the quantitative calculation of solvation phenomena. *J. Phys. Chem. A.* **1995**, *95*, 2224–2235.
3. Preiss. P.R.M.U.; Slattery. M.J.; Krossing. I. In silico prediction of molecular volumes, heat capacities and temperature dependent

- densities of ionic liquids. *Ind. Eng. Chem. Res.* **2009**, *48*, 2290–2296.
4. Diedenhofen, M.; Klamt, A. COSMO-RS as a tool for property prediction of IL mixture – A review. *Fluid Phase Equil.* **2010**, *294*, 31–38.