



národní  
úložiště  
šedé  
literatury

### **A Note on Dempster Rule**

Hájek, Petr  
2003

Dostupný z <http://www.nusl.cz/ntk/nusl-34140>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 24.09.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní [nusl.cz](http://nusl.cz).



**Institute of Computer Science**  
**Academy of Sciences of the Czech Republic**

## **A NOTE ON DEMPSTER RULE**

PETR HÁJEK and DAGMAR HARMANCOVÁ

Technical report No. 891

January 2003



**Institute of Computer Science**  
**Academy of Sciences of the Czech Republic**

## **A NOTE ON DEMPSTER RULE**

PETR HÁJEK and DAGMAR HARMANCOVÁ<sup>1</sup>

Technical report No. 891

January 2003

Abstract:

Dempster rule of combination of belief functions is shown not to commute with restriction to a sublanguage – badly for one version of the rule and less badly, but still for an alternative version.

Keywords:

Dempster-Shafer theory, belief functions, Dempster rule

---

<sup>1</sup>Institute of Computer Science, Academy of Sciences of the Czech Republic, Pod Vodárenskou věží 2, 182 07 Prague, Czech Republic, Phone: +420 266 053 760

For basic information on Dempster-Shafer theory (belief function, basic belief [probability] assignment] etc.) see [1]. Note that Shafer deals only with regular belief functions, i.e. assigning 0 to empty set. This is generalized in [2] (see also [3]). Here we deal with general (both regular and singular) belief functions. If  $m$  is basic belief assignment then the corresponding belief function is  $bel(A) = \sum_{B \subseteq A} m(B)$ . Note that  $bel(A) + bel(W - A) \leq 1 + m(\emptyset)$ . Dempster rule is a famous operation assigning to two belief functions  $bel_1, bel_2$  on the same (finite) set  $W$  their combination  $bel_1 \oplus bel_2$ . It can be defined using the corresponding basic belief assignments  $m_1, m_2$ , defining, for  $A \subseteq W$ ,  $(bel_1 \oplus bel_2)(A) = \sum_{B \cap C \subseteq A} m_1(B) \cdot m_2(C)$ . (Note that  $bel_1 \oplus bel_2$  need not be regular even if both  $bel_1$  and  $bel_2$  is.) Alternatively, one can use Dempster spaces (as defined in the pioneering paper [4])  $\mathbf{D}_i = (E_i, W, \Gamma_i, \mu_i) (i = 1, 2)$  and define their product to be  $\mathbf{D} = (E, W, \Gamma, \mu)$  where  $E = E_1 \times E_2$ ,  $\Gamma(e_1, e_2) = \Gamma_1(e_1) \cap \Gamma_2(e_2)$  and  $\mu$  is the product measure ( $\mu(e_1, e_2) = \mu_1(e_1) \cdot \mu_2(e_2)$ ). If  $bel_i$  is given by  $\mathbf{D}_i$  then  $bel_1 \oplus bel_2$  is given by  $\mathbf{D}$ .

When we are interested in belief functions on (classes of equivalent boolean) formulas and consider formulas built from finitely many variables  $p_1, \dots, p_n$  then we may identify formulas with subsets of  $2^n$  (sets of  $n$ -tuples of zeros and ones) in the obvious way; then we may compute, given  $bel_i$ , the corresponding belief assignments and define  $bel_1 \oplus bel_2$ .

Let  $B_n$  be the algebra of formulas built from propositional variables  $p_1, \dots, p_n$ . If  $k < n$  then  $B_k$  is a subalgebra of  $B_n$  and the restriction of a belief function on  $B_n$  to  $B_k$  is a belief function in  $B_k$ . This is immediate using the condition of superadditivity: A function  $bel$  mapping the algebra of (classes of equivalent) formulas into  $[0, 1]$  is a belief function (see [1]) iff  $bel(true) = 1$  and

$$bel(\varphi_1 \vee \dots \vee \varphi_n) \geq \sum_{\emptyset \neq I \subseteq \{1, \dots, n\}} (-1)^{|I|+1} bel\left(\bigwedge_{i \in I} \varphi_i\right).$$

A binary operation  $F$  on belief functions (on the same algebra) *commutes with restriction* if for any pair  $bel_1, bel_2$  of belief functions on  $B_n$  and for  $k < n$ ,

$$F(bel_1, bel_2) \upharpoonright B_k = F(bel_1 \upharpoonright B_k, bel_2 \upharpoonright B_k)$$

A trivial example is a convex combination: for  $0 < \alpha, \beta < 1$  and  $\alpha + \beta = 1$ ,  $F(bel_1, bel_2)(A) = \alpha bel_1(A) + \beta bel_2(A)$ .

Our question is, if Dempster rule commutes with restriction, i.e. if we consider formulas from  $B_7$ , say, i.e. built using  $p_1, \dots, p_7$  and among them we take a  $\varphi$  containing only  $p_1, p_2, p_3$ , can we compute  $(bel_1 \oplus bel_2)(\varphi)$  working only with restriction of  $bel_i$  to  $B_3$ ? The answer is NO – for our definition of  $bel$ . The following is an example, with  $n = 2$  and  $k = 1$ ,  $bel_1 = bel_2$ ,  $\varphi$  is  $p$ ,  $bel_1(p) = 0.9$ ,  $m'_1 = m'_2$  is the *b.b.a* for  $bel_1 \upharpoonright B_1$ . Furthermore,  $bel = bel_1 \oplus bel_2$ ,  $bel' = bel'_1 \oplus bel'_2$ .

	$p \& q$	$p \& \neg q$	true
$m_1$	.2	.7	.1

  

	$p$	true
$m'_1$	0.9	0.1

  

	$p \& q$	$p \& \neg q$	true
$p \& q$	$p \& q$	false	$p \& q$
	.04	.14	.02
$p \& \neg q$	false	$p \& \neg q$	$p \& \neg q$
	.14	.49	.07
true	$p \& q$	$p \& \neg q$	true
	.02	.07	.01

  

	$p$	true
$p$	$p$	$p$
	.81	.09
true	$p$	true
	.09	.01

	false	$p \& q$	$p \& \neg q$	true
$m_1 \oplus m_2$	.28	.08	.63	.01

	$p$	true
$m'_1 \oplus m'_2$	.99	.01

Observe the following:  $bel(p) = bel'(p) = 0.99$ ,  $bel(\neg p) = bel(false) = 0.28$ ,  $bel'(\neg p) = bel'(false) = 0$ .

Let us consider restriction from  $B_n$  to  $B_{n-1}$ , or, equivalently, the belief function  $bel'$  on subsets of  $2^{n-1}$  induced by  $bel$  on subsets of  $2^n$ . Let  $A, B, C$  run on subsets of  $2^{n-1}$  and  $U, V, W$  on subsets of  $2^n$ . Define

$$ext(A) = \{ \langle \varepsilon_1, \dots, \varepsilon_n \rangle \mid \langle \varepsilon_1, \dots, \varepsilon_{n-1} \rangle \in A, \varepsilon_n \in \{0, 1\} \},$$

$$proj(U) = \{ \langle \varepsilon_1, \dots, \varepsilon_{n-1} \rangle \mid \text{for some } \varepsilon_n, \langle \varepsilon_1, \dots, \varepsilon_{n-1}, \varepsilon_n \rangle \in U \},$$

(extension and projection). Clearly,  $bel'(A) = bel(ext(A))$ . If  $m$  is the b.b.a. of  $bel$ , put

$$m'(A) = \sum_{proj(U)=A} m(U);$$

this is the b.b.a. of  $bel'$ . Indeed, clearly  $\sum_{A \subseteq 2^{n-1}} m'(A) = 1$ ; moreover,

$$\begin{aligned} bel'(A) &= bel(ext(A)) = \sum_{U \subseteq ext(A)} m(U) = \sum_{proj(U) \subseteq A} m(U) = \\ &= \sum_{B \subseteq A} \sum_{proj(U)=B} m(U) = \sum_{B \subseteq A} m'(B). \end{aligned}$$

### Lemma

- (1) Under the above notation,  $proj(U \cap V) \subseteq proj(U) \cap proj(V)$ ; but there are  $U, V$  for which this inclusion is proper.
- (2) For each  $A \subseteq 2^{n-1}$ ,  $proj(U) \subseteq A$  iff  $U \subseteq ext(A)$ .

### Proof.

- (1) Evidently  $(U \cap V) \subseteq U$  and hence  $proj(U \cap V) \subseteq proj(U)$ , similarly  $proj(U \cap V) \subseteq proj(V)$  and hence  $proj(U \cap V) \subseteq proj(U) \cap proj(V)$ . For a counterexample take  $n = 2$ ,  $U = \{(1, 0)\}$ ,  $V = \{(1, 1)\}$ ; then  $U \cap V = \emptyset = proj(U \cap V)$ , but  $proj(U) = proj(V) = \{(1)\} = proj(U) \cap proj(V)$ .
- (2) is obvious. □

**Theorem 1.** Let, for  $i = 1, 2$   $bel_i$  be a belief function on  $B_n$  and  $bel'_i$  its restriction to  $B_{n-1}$ . Let  $bel = bel_1 \oplus bel_2$ ,  $bel' = bel'_1 \oplus bel'_2$ . Then, for each  $\varphi \in B_{n-1}$ ,

$$bel'(\varphi) \leq bel(\varphi)$$

.

**Proof.** Let  $A$  be the set of  $(n-1)$ -tuples satisfying  $\varphi$ ; we claim  $bel'(A) \leq bel(ext(A))$ .

We compute:

$$\begin{aligned} (bel'_1 \oplus bel'_2)(A) &= \sum_{B \cap C \subseteq A} m'_1(B) \cdot m'_2(C) = \\ &= \sum_{B \cap C \subseteq A} \left( \sum_{proj(U)=B} m_1(U) \cdot \sum_{proj(V)=C} m_2(V) \right) = \\ &= \sum_{proj(U) \cap proj(V) \subseteq A} m_1(U) \cdot m_2(V) \leq \sum_{proj(U \cap V) \subseteq A} m_1(U) \cdot m_2(V) = \end{aligned}$$

$$\begin{aligned}
&= \sum_{U \cap V \subseteq \text{ext}(A)} m_1(U) \cdot m_2(V) = \\
&= \text{bel}_1 \oplus \text{bel}_2(\text{ext}(A))
\end{aligned}$$

□

\*

So far so good; but now consider another (and more usual) definition of the belief function given by a possibly singular b.b.a.  $m$ , namely  $\text{Bel}(A) = \sum_{\emptyset \neq B \subseteq A} m(B)$  (this works for  $A \neq \emptyset$ ; one puts  $\text{Bel}(\emptyset) = 0$ ). This may be not-normalized ( $\text{Bel}(\text{true})$  may be  $< 1$ ). A normalized belief function is then  $n\text{Bel}(A) = (\sum_{\emptyset \neq B \subseteq A} m(B)) / (\sum_{B \neq \emptyset} m(B))$ . Dempster rule for (normalized) belief functions  $\text{Bel}_1, \text{Bel}_2$  is then defined for  $A \neq \emptyset$  as

$$(\text{Bel}_1 \oplus \text{Bel}_2)(A) = \sum_{\emptyset \neq B \cap C \subseteq A} m_1(B) \cdot m_2(C),$$

(non-empty subsets!),  $(\text{Bel}_1 \oplus \text{Bel}_2)(\emptyset) = 0$ . This may give a non-normalized belief function; it can be normalized by dividing by  $1 - \sum_{B \cap C = \emptyset} m_1(B) \cdot m_2(C)$  (if this is non-zero). Even if reasonably motivated, it has the following (unwanted?) property:

**Observation.** Let  $m_i, \text{Bel}_i, \text{Bel}'_i, \text{Bel}, \text{Bel}'$  be as above but belief functions in the new meaning just defined; Dempster rule either normalized or not. Then no inequality can be proved for  $\text{Bel}(A), \text{Bel}'(A)$ ; all three possibilities

$$\text{Bel}'(A) < \text{Bel}(A), \text{Bel}'(A) = \text{Bel}(A), \text{Bel}'(A) > \text{Bel}(A)$$

may occur (and similarly for  $n\text{Bel}', n\text{Bel}$ ).

For  $\text{Bel}'(p) > \text{Bel}(p)$  consider the example above:  $(\text{Bel}_1 \oplus \text{Bel}_1)(p) = .08 + .63 = .71$  (normalized:  $71/72$ ) but  $(\text{Bel}'_1 \oplus \text{Bel}'_2)(p) = .99$  (normal).

Now  $\text{Bel}'(\varphi) = \text{Bel}(\varphi)$  as well as  $\text{bel}'(\varphi) = \text{bel}(\varphi)$  holds for all  $\varphi \in B_{n-1}$  if all focal elements have the form  $\text{ext}(A)$  for some  $A \subseteq 2^n$  (verify!). In general, if  $m$  is a b.b.a. and  $\text{bel}$  is the corresponding belief function in our former sense then  $\text{Bel}(\varphi) = \text{bel}(\varphi) - \text{bel}(\text{false})$  is the (possibly non-normalized) belief function in the latter sense and  $n\text{Bel}(\varphi) = (\text{bel}(\varphi) - \text{bel}(\text{false})) / (1 - \text{bel}(\text{false}))$  is its normalisation. Now if  $\text{bel} = \text{bel}_1 \oplus \text{bel}_2$  and  $\text{bel}' = \text{bel}'_1 \oplus \text{bel}'_2$  in the former sense then Dempster sum in the latter sense of  $\text{Bel}_1, \text{Bel}_2$  is  $\text{bel} - \text{bel}(\text{false})$  and its normalized  $n\text{Bel}$  is  $(\text{bel} - \text{bel}(\text{false})) / (1 - \text{bel}(\text{false}))$ ; similarly for  $\text{bel}'_i, \text{bel}'$ . Put  $x = \text{bel}'(\varphi), y = \text{bel}(\varphi), c = \text{bel}'(\text{false}), d = \text{bel}(\text{false})$ . Then  $x \leq y$  and  $c \leq d$  (by Theorem 1) and  $\text{Bel}'(\varphi) \leq \text{Bel}(\varphi)$  iff  $x - c \leq y - d$  iff  $(d - c) \leq (y - x)$  and  $n\text{Bel}'(\varphi) \leq n\text{Bel}(\varphi)$  iff  $\frac{x-c}{1-c} \leq \frac{y-d}{1-d}$  iff  $(d - c) \leq y(1 - c) - x(1 - d)$  (by elementary computations). In particular, the following is an example for  $\text{Bel}'(\varphi) < \text{Bel}(\varphi)$  and for  $n\text{Bel}'(\varphi) < n\text{Bel}(\varphi)$ :

$$\begin{array}{lll}
m_1(q) = 0.5 & m_1(\text{true}) = 0.5 & m'_1(p) = m'_2(p) = 0 \\
m_2(p \equiv q) = 0.5 & m_2(\text{true}) = 0.5 & m'_1(\text{true}) = m'_2(\text{true}) = 0
\end{array}$$

then

$$\text{bel}(p) = 0.25 = \text{Bel}(p) = n\text{Bel}(p) \quad \text{bel}'(p) = 0 = \text{Bel}'(p) = n\text{Bel}'(p)$$

## 1 Conclusion.

As commonly known, the set  $BEL(B_n)$  all belief functions on algebra  $B_n$  endowed with the operation  $\oplus$  makes  $BEL$  to a commutative semigroup with a unit element. The natural projection of  $B_n$  onto  $B_{n-1}$  (or to  $B_k, k < n$ ) induces a natural projection of  $BEL(B_n)$  onto  $BEL(B_{n-1})$ . The fact that this projection does not commute with the operation  $\oplus$  (i.e. it is not a semigroup homomorphism) may be disappointing; but, to take it positively, contributes to our understanding of Dempster rule, which has proven to be useful in many situation. From possible variants of the definition of Dempster

rule, our first variant shows to have smoother properties than the other ones; but on the other hand, the fact that  $bel(\varphi) + bel(\neg\varphi)$  may be more than 1 may be felt to be counter-intuitive (if not carefully interpreted). One can ask if there is still other natural variant of Dempster-style combination of belief functions that would commute with taking subalgebra. This remains as an interesting open problem.

**Remark.** In this context let us mention that in [5] (section 2.4) the authors consider a related but different problem that can be formulated using our framework as follows: Let  $B_n$  be given and let  $B_{n_1}, B_{n_2}$  be subalgebra of formulas built from propositional variables  $p_1, \dots, p_i$  and  $p_{i+1}, \dots, p_n$  respectively. Let  $Bel$  be a normalized belief function on  $B_n$  and let  $Bel_1, Bel_2$  be projections to  $B_{n_1}, B_{n_2}$  respectively; let  $Bel_1', Bel_2'$  be their extensions to  $B_n$  and let  $Bel^* = Bel_1' \oplus Bel_2'$ . Can one conclude  $Bel(A) \leq Bel^*(A)$  or  $Bel^*(A) \leq Bel(A)$ ? They give examples showing that the answer is *no*. (Thanks are due to D. Dubois for calling our attention to [5].)

### Acknowledgement

Partial support by the grant No. AA1030004/00 of the Grant Agency of the Academy of Sciences of the Czech Republic is acknowledged.

## Bibliography

- [1] G. Shafer, *A Mathematical Theory of Evidence* (Princeton Univ. Press 1975).
- [2] Ph. Smets, “The combination of evidence in transferable belief model”, *IEEE Trans. Pattern Anal. and Machine Int.* **12** (1990) 447–458.
- [3] P. Hájek, “Getting belief functions from Kripke models”, *Int. J. General Systems*, **24** (1996) 325–327
- [4] A. P. Dempster, “Upper and lower probabilities induced by a multi-valued mapping”, *Ann. Math. Stat.* **38** (1967) 325–ff.
- [5] D. Dubois and H. Prade, “A set-theoretic view of belief functions”, *Int. J. General Systems* **12** (1986) 193–226.