



národní  
úložiště  
šedé  
literatury

**Regional Air Pollution Transport and Its Influence to Pollution Hotspot in the Czech Republic.**

Kozáková, Jana  
2017

Dostupný z <http://www.nusl.cz/ntk/nusl-369613>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 18.04.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní [nusl.cz](http://nusl.cz).

## REGIONAL AIR POLLUTION TRANSPORT AND ITS INFLUENCE TO POLLUTION HOTSPOT IN THE CZECH REPUBLIC

Jana Kozáková<sup>1,2</sup>, Petr Vodička<sup>1</sup>, Petra Pokorná<sup>1,2</sup>, Lucie Ondráčková<sup>1</sup>, Jakub Ondráček<sup>1</sup>, Jan Hovorka<sup>2</sup>, Kamil Křůmal<sup>3</sup>, Pavel Mikuška<sup>3</sup>, Pavel Moravec<sup>1</sup>, Jaroslav Schwarz<sup>1</sup>

<sup>1</sup>Institute of Chemical Process Fundamentals of the CAS, v.v.i., Laboratory of Aerosols Chemistry and Physics, Rozvojová 135, Prague 165 02, Czech Republic

<sup>2</sup>Charles University, Faculty of Science, Institute for Environmental Studies, Benátská 2, Prague 128 01, Czech Republic; kozakova@icpf.cas.cz

<sup>3</sup>Institute of Analytical Chemistry of the CAS, v.v.i., Veveří 97, 602 00 Brno, Czech Rep.

Keywords: Chemical composition, PM<sub>1</sub>, PM<sub>10</sub>, Urban aerosol

### INTRODUCTION

Upper Silesian Industrial Region (USIR) represents large area with enormous concentration of industry and poor air quality. Dominant part of USIR is located in the Polish Silesian Province and small part in Moravian-Silesian Region in the Czech Republic where one of the European air pollution hotspots is the city Ostrava (Pokorná et al., 2015). Elevated air pollution level has been associated with cardiovascular and respiratory diseases in adults and children and other health consequences (WHO, 2013).

The aim of the study was to find out the local effect and the regional air pollution transport effect on the air quality of the Ostrava Radvanice district, the problematic air pollution hot spot.

### EXPERIMENTAL SETUP

Sampling sites were located at urban industrial site Ostrava-Radvanice a Bartovice and suburban site Ostrava-Plesná in the winter period 6.2.2014-6.3.2014.

24h mass concentrations of PM<sub>10</sub> and PM<sub>1</sub> were determined and size resolved mass concentrations of atmospheric aerosol (AA) were measured by cascade impactors. Chemical analyses of PM<sub>10</sub>, PM<sub>1</sub>, and size resolved AA were performed. Meteorological parameters from nearest monitoring stations were used to determine the association between air pollution level and meteorological situation.

### RESULTS AND CONCLUSIONS

Higher average mass concentrations of PM and chemical species were measured in Radvanice than in Plesná, except NO<sub>3</sub><sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, and NH<sub>4</sub><sup>+</sup>, in both PM fractions. Na<sup>+</sup> and Ca<sup>2+</sup> in PM<sub>1</sub> were four times higher and Ca<sup>2+</sup> and Mg<sup>2+</sup> in PM<sub>10</sub> even fifth times higher in Radvanice than in Plesná (Table 1). It suggests that Radvanice site was influenced by a close industrial zone (IZ) located in the southwest (SW) direction from measurement station.

To investigate the influence of the IZ in Radvanice, ratios between mass concentrations measured in Plesná and Radvanice were calculated (Figure 1) during SW and northeast (NE) wind directions (WDs). During SW WD mass concentrations were higher in Radvanice than in Plesná and thus the influence of the IZ was evident. This

finding was also confirmed by increased mass concentrations of specific chemical species which are generally produced by the industrial sources. The effect of the regional air pollution transport was evident in Plesná during NE WD when the mass concentrations of the most of species were increased.

Tab. 1: Average 24h mass concentrations and STD for selected species.

(µg.m <sup>-3</sup> )	Radvanice		Plesná	
	PM <sub>1</sub>	PM <sub>10</sub>	PM <sub>1</sub>	PM <sub>10</sub>
PM	29.4±9.2	55.0±18.0	28.8±15.5	39.6±22.4
OC	11.3±8.5	14.6±13.6	7.0±4.3	9.7±6.6
EC	2.1±1.8	3.3±3.6	2.3±1.3	2.5±1.6
NO <sub>3</sub> <sup>-</sup>	2.2±1.0	4.1±1.9	3.1±1.3	4.6±2.0
SO <sub>4</sub> <sup>2-</sup>	2.3±1.1	4.3±2.4	3.2±2.0	4.6±3.2
NH <sub>4</sub> <sup>+</sup>	1.5±0.7	2.1±1.4	2.3±1.2	3.0±1.8
(ng.m <sup>-3</sup> )				
Ca <sup>2+</sup>	66±43	780±554	15±9	155±123
Na <sup>+</sup>	267±222	468±180	61±66	212±118
Mg <sup>2+</sup>	6±8	123±103	2±1	27±16
Fe	261±262	-	83±95	-




Fig. 1: Ratios between mass concentrations at Plesná and Radvanice sites.

#### ACKNOWLEDGEMENT

This work was supported by the Czech Grant Agency (P503/12/G147).

#### REFERENCES

- Pokorná, P., Hovorka, J., Klán, M., Hopke, P.K., Source Apportionment of Size Resolved Particulate Matter at a European Air Pollution Hot Spot, *Sci. Total Environ.*, 502, 172–183, (2015).
- WHO, 2013. Review of Evidence on Health Aspects of Air Pollution – REVIHAAP Project, WHO.