



národní
úložiště
šedé
literatury

On Rump's Characterization of P-Matrices

Rohn, Jiří
2010

Dostupný z <http://www.nusl.cz/ntk/nusl-41912>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 08.05.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní [nusl.cz](http://www.nusl.cz).



Institute of Computer Science
Academy of Sciences of the Czech Republic

On Rump's Characterization of *P*-Matrices

Jiří Rohn

Technical report No. V-1093

30.11.2010



Institute of Computer Science
Academy of Sciences of the Czech Republic

On Rump's Characterization of *P*-Matrices

Jiří Rohn¹

Technical report No. V-1093

30.11.2010

Abstract:

Necessary and sufficient *P*-matrix condition by S. M. Rump is simplified by showing that one of its assumptions can be deleted without affecting validity of the result.

Keywords:

P-matrix, interval matrix, regularity.

¹This work was supported by the Czech Republic Grant Agency under grants 201/09/1957 and 201/08/J020, and by the Institutional Research Plan AV0Z10300504.

1 Introduction

S. M. Rump in his paper [2] proved the following result concerning a square matrix A .

Theorem 1. *Let both $A - I$ and $A + I$ be nonsingular. Then A is a P -matrix if and only if the interval matrix*

$$[(A - I)^{-1}(A + I) - I, (A - I)^{-1}(A + I) + I] \quad (1.1)$$

is regular.

Here, I is the identity matrix and the interval matrix (1.1) is called regular if each point matrix contained therein is nonsingular. The main purpose of the present paper consists in showing that the assumption of nonsingularity of $A + I$ is redundant and may be deleted without affecting validity of the theorem. The main result (Theorem 4) is preceded by two auxiliary propositions that make it easier to understand the relationship between the P -property of real matrices and regularity of interval matrices.

2 Definitions and notations

Given an $n \times n$ matrix A and a subset $\emptyset \neq J \subseteq \{1, \dots, n\}$, denote by $A[J]$ the submatrix of A consisting of rows and columns whose indices belong to J . Submatrices formed in this way are called principal submatrices, and A is said to be a P -matrix if determinants of all the principal submatrices (also called principal minors) are positive.

We have defined, as usual, $A[J]$ for $J \neq \emptyset$ only, but we shall also need to have it defined for $J = \emptyset$. In this case we define $A[\emptyset]$ to be the empty matrix, and we set $\det(A[\emptyset]) = 1$.

For each vector $y \in Y_n$ we put

$$A_y = (A - I)^{-1}(A + I) - T_y, \quad (2.1)$$

where $T_y = \text{diag}(y)$ is the diagonal matrix with diagonal vector y . Obviously,

$$A_y \in [(A - I)^{-1}(A + I) - I, (A - I)^{-1}(A + I) + I]$$

for each such a y .

3 The result

First we show that regularity of the interval matrix (1.1) can be formulated in terms of determinants of the matrices A_y .

Proposition 2 *Let $A - I$ be nonsingular. Then the interval matrix*

$$[(A - I)^{-1}(A + I) - I, (A - I)^{-1}(A + I) + I]$$

is regular if and only if the numbers

$$\det(A_y), \quad y \in Y_n$$

are either all simultaneously positive, or all simultaneously negative.

Proof. In [1, Thm. 5.1, (C1)] it is proved that an interval matrix $[A_c - \Delta, A_c + \Delta]$ is regular if and only if the determinants of the matrices

$$A_c - T_{y'} \Delta T_{z'}, \quad y', z' \in Y_n$$

are either all simultaneously positive, or all simultaneously negative. In our case we have

$$A_c - T_{y'} \Delta T_{z'} = (A - I)^{-1}(A + I) - T_y = A_y$$

where y is given by $y_i = y'_i z'_i$ ($i = 1, \dots, n$), hence the general condition reduces to the respective property of matrices A_y , $y \in Y_n$. \square

Next we show a connection between determinants of matrices A_y and principal minors of A .

Proposition 3 *Let $A - I$ be nonsingular. Then for each $y \in Y_n$ we have*

$$\det(A_y) = \frac{2^n \det(A[J(y)])}{\det(A - I)}, \quad (3.1)$$

where

$$J(y) = \{j \mid y_j = -1\}.$$

Proof. Let $y \in Y_n$. Then from (2.1) we have

$$(A - I)A_y = A + I - (A - I)T_y = A(I - T_y) + I + T_y \quad (3.2)$$

and for the j th column of the right-hand side matrix there holds

$$(A(I - T_y) + I + T_y)_{\bullet j} = \begin{cases} 2I_{\bullet j} & \text{if } y_j = 1, \\ 2A_{\bullet j} & \text{if } y_j = -1 \end{cases}$$

($j = 1, \dots, n$). Taking the Laplace expansion along all the columns with $y_j = 1$, we obtain

$$\det(A(I - T_y) + I + T_y) = 2^n \det(A[J(y)]) \quad (3.3)$$

where

$$J(y) = \{j \mid y_j = -1\},$$

and this result also holds for $J(y) = \emptyset$ because in this case both sides in (3.3) are equal to 2^n in view of our definition of $\det(A[\emptyset])$ in Section 2. Hence, from (3.2) and (3.3) we obtain

$$\det(A - I) \det(A_y) = 2^n \det(A[J(y)])$$

and

$$\det(A_y) = \frac{2^n \det(A[J(y)])}{\det(A - I)},$$

which concludes the proof. \square

Finally, we prove our main result.

Theorem 4. *Let $A - I$ be nonsingular. Then A is a P -matrix if and only if the interval matrix*

$$[(A - I)^{-1}(A + I) - I, (A - I)^{-1}(A + I) + I] \quad (3.4)$$

is regular.

Proof. Let $A - I$ be nonsingular. If A is a P -matrix, then $\det(A[J(y)]) > 0$ for each $y \in Y_n$ and by (3.1) the determinants of all the matrices A_y , $y \in Y_n$ are of the same sign (positive if $\det(A - I) > 0$ and negative if $\det(A - I) < 0$), hence the interval matrix (3.4) is regular due to Proposition 2. Conversely, if the interval matrix (3.4) is regular, then determinants of all the matrices A_y are of the same sign by Proposition 2 which in view of (3.1) means that all the numbers $\det(A[J(y)])$, $y \in Y_n$ are of the same sign. But since for $y = (1, \dots, 1)^T$ we have $\det(A[J(y)]) = \det(A[\emptyset]) = 1$, all the principal minors are positive and A is a P -matrix. \square

This proves our original claim.

Bibliography

- [1] J. Rohn, *Systems of linear interval equations*, Linear Algebra and Its Applications, 126 (1989), pp. 39–78. [3](#)
- [2] S. M. Rump, *On P-matrices*, Linear Algebra and Its Applications, 363 (2003), pp. 237–250. [2](#)