



národní
úložiště
šedé
literatury

A Perturbation Theorem for Linear Equations

Rohn, Jiří
2011

Dostupný z <http://www.nusl.cz/ntk/nusl-42443>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 30.09.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní [nusl.cz](http://www.nusl.cz) .



Institute of Computer Science
Academy of Sciences of the Czech Republic

A Perturbation Theorem for Linear Equations

Jiří Rohn

Technical report No. V-1103

18.01.2011



Institute of Computer Science
Academy of Sciences of the Czech Republic

A Perturbation Theorem for Linear Equations

Jiří Rohn¹

Technical report No. V-1103

18.01.2011

Abstract:

This is an unpublished two-page manuscript from 2000. We describe explicit formulae for componentwise bounds on solution of a system of linear equations $A_c x = b_c$ (A_c square) under perturbation of all data. To make the result numerically tractable, we avoid use of exact inverses, using instead some matrices R and M required only to satisfy certain inequalities. Hansen's optimality result is a special case of our theorem.

Keywords:

Linear equations, perturbation, bounds.

¹This work was supported by the Institutional Research Plan AV0Z10300504.

ROHN, J.

A perturbation theorem for linear equations

We describe explicit formulae for componentwise bounds on solution of a system of linear equations $A_c x = b_c$ (A_c square) under perturbation of all data. To make the result numerically tractable, we avoid use of exact inverses, using instead some matrices R and M required only to satisfy certain inequalities. Hansen's optimality result is a special case of our theorem.

Notations used: I is the unit matrix, ϱ denotes the spectral radius, for $A = (a_{ij})$ we denote $|A| = (|a_{ij}|)$ and inequalities are understood componentwise. To save space, we write a/b instead of $\frac{a}{b}$.

Theorem. Let $M \geq 0$ and R be arbitrary matrices satisfying

$$MG + I \leq M, \quad (1)$$

where $G = |I - RA_c| + |R|\Delta$. Then for each A and b such that $|A - A_c| \leq \Delta$ and $|b - b_c| \leq \delta$, A is nonsingular and the solution of the system $Ax = b$ satisfies for each $i \in \{1, \dots, n\}$

$$\min\{\underline{x}_i/\alpha_i, \underline{x}_i/\beta_i\} \leq x_i \leq \max\{\tilde{x}_i/\alpha_i, \tilde{x}_i/\beta_i\}, \quad (2)$$

where

$$\begin{aligned} \underline{x}_i &= -(M(|Rb_c| + |R|\delta))_i + m_i(Rb_c + |Rb_c|)_i \\ \tilde{x}_i &= (M(|Rb_c| + |R|\delta))_i + m_i(Rb_c - |Rb_c|)_i \\ \alpha_i &= 1 + (|r_i| - r_i)m_i + h_i \\ \beta_i &= 2m_i - 1 - (|r_i| + r_i)m_i - h_i \\ m_i &= M_{ii} \\ r_i &= (I - RA_c)_{ii} \\ h_i &= (M - MG - I)_{ii} \end{aligned}$$

and $\beta_i \geq \alpha_i \geq 1$. Moreover, if $A_c = I$ and $\varrho(\Delta) < 1$ and if we take $R = I$ and $M = (I - \Delta)^{-1}$, then the bounds (2) are exact (i.e., attained).

Proof. 1) First we prove that each matrix A with $|A - A_c| \leq \Delta$ is nonsingular. Premultiplying the inequality (1) by the nonnegative matrix G yields $MG^2 + G + I \leq MG + I \leq M$ and by induction $\sum_{j=0}^k G^j \leq MG^{k+1} + \sum_{j=0}^k G^j \leq M$ for each $k \geq 0$, hence $\sum_{j=0}^{\infty} G^j$ is convergent which, as well known, implies that $\varrho(G) < 1$. Since $|I - RA| = |I - RA_c + R(A_c - A)| \leq |I - RA_c| + |R|\Delta = G$, we have $\varrho(I - RA) \leq \varrho(G) < 1$ which means that the matrix $RA = I - (I - RA)$ is nonsingular, hence A is nonsingular.

2) Next we prove that $\beta_i \geq \alpha_i \geq 1$ for each i . From the definition of h_i we have $m_i = (MG)_{ii} + 1 + h_i \geq m_i|r_i| + 1 + h_i$ which can be easily rearranged to $2m_i - 1 - (|r_i| + r_i)m_i - h_i \geq 1 + (|r_i| - r_i)m_i + h_i$, giving $\beta_i \geq \alpha_i$; the inequality $\alpha_i \geq 1$ follows from the nonnegativity of m_i and h_i .

3) Let x solve $Ax = b$ for some A, b with $|A - A_c| \leq \Delta$ and $|b - b_c| \leq \delta$. Then we have

$$x = (I - RA)x + Rb = (I - RA_c)x + R(A_c - A)x + Rb_c + R(b - b_c) \quad (3)$$

and taking absolute values gives

$$|x| \leq G|x| + |Rb_c| + |R|\delta. \quad (4)$$

Let $i \in \{1, \dots, n\}$. Then from the i th equation in (3) we have

$$\begin{aligned} x_i &\leq ((I - RA_c)x)_i + (|R|\Delta|x|)_i + (Rb_c)_i + (|R|\delta)_i \\ &= (G|x| + |Rb_c| + |R|\delta)_i + ((I - RA_c)x - |I - RA_c| \cdot |x| + Rb_c - |Rb_c|)_i. \end{aligned} \quad (5)$$

Put $x' = (|x_1|, \dots, |x_{i-1}|, x_i, |x_{i+1}|, \dots, |x_n|)^T$. Then from (4) and (5) we have $x' \leq G|x| + |Rb_c| + |R|\delta + ((I - RA_c)x - |I - RA_c| \cdot |x| + Rb_c - |Rb_c|)_i e_i$, where e_i is the i th column of I . Premultiplying this inequality by the nonnegative vector $e_i^T M$ and using the matrix $H := M - MG - I$, we obtain $(Mx')_i \leq ((M - H - I)|x|)_i + ((I - RA_c)x - |I - RA_c| \cdot |x|)_i m_i + \tilde{x}_i$ and consequently

$$(M(x' - |x|))_i + (H|x|)_i + |x_i| + (|I - RA_c| \cdot |x| - (I - RA_c)x)_i m_i \leq \tilde{x}_i. \quad (6)$$

Since $(M(x' - |x|))_i = m_i(x_i - |x_i|)$, $(H|x|)_i \geq h_i|x_i|$ and $(|I - RA_c| \cdot |x| - (I - RA_c)x)_i \geq |r_i| \cdot |x_i| - r_i x_i$, from (6) we finally obtain an inequality containing variable x_i only:

$$m_i(x_i - |x_i|) + h_i|x_i| + |x_i| + (|r_i| \cdot |x_i| - r_i x_i)m_i \leq \tilde{x}_i. \quad (7)$$

If $x_i \geq 0$, then this inequality becomes $\alpha_i x_i \leq \tilde{x}_i$, implying $x_i \leq \tilde{x}_i/\alpha_i$, and if $x_i < 0$, then (7) turns into $\beta_i x_i \leq \tilde{x}_i$, giving $x_i \leq \tilde{x}_i/\beta_i$, which together yields

$$x_i \leq \max\{\tilde{x}_i/\alpha_i, \tilde{x}_i/\beta_i\}. \quad (8)$$

In this way we have obtained the upper bound in (2). To prove the lower one, notice that $-x$ satisfies $A(-x) = -b$, where $|A - A_c| \leq \Delta$ and $|(-b) - (-b_c)| \leq \delta$. Hence we can use the previously obtained result if we set $b_c := -b_c$, which affects \tilde{x}_i only. Then from (8) we get $-x_i \leq \max\{-\tilde{x}_i/\alpha_i, -\tilde{x}_i/\beta_i\}$ which, after premultiplying by -1 , gives the lower bound in (2).

4) Finally, to prove the optimality result for the case $A_c = I$ and $\varrho(\Delta) < 1$, take $R = I$ and $M = (I - \Delta)^{-1}$, then $M \geq 0$, $G = \Delta$ and (1) is satisfied as an equation; moreover, for each i we have $r_i = h_i = 0$, $\alpha_i = 1$, $\beta_i = 2m_i - 1$, hence (2) has the form

$$\min\{\tilde{x}_i, \tilde{x}_i/\beta_i\} \leq x_i \leq \max\{\tilde{x}_i, \tilde{x}_i/\beta_i\}. \quad (9)$$

To prove that the upper bound is really attained, let us fix an $i \in \{1, \dots, n\}$ and define a diagonal matrix D by $D_{jj} = 1$ if $j \neq i$ and $(b_c)_j \geq 0$, $D_{jj} = -1$ if $j \neq i$ and $(b_c)_j < 0$, and $D_{jj} = 1$ if $j = i$, and let $\tilde{b} = Db_c + \delta$. Then it can be easily verified that $\tilde{x}_i = (M\tilde{b})_i$ holds. First, define $x' = DM\tilde{b}$. Since $M = (I - \Delta)^{-1}$ implies $\Delta M = M\Delta = M - I$, we have $(I - D\Delta D)x' = DM\tilde{b} - D(M - I)\tilde{b} = D\tilde{b} = b_c + D\delta$, which means that x' solves the system $(I - D\Delta D)x' = b_c + D\delta$ where $|(I - D\Delta D) - I| = \Delta$, $|(b_c + D\delta) - b_c| = \delta$ and $x'_i = e_i^T DM\tilde{b} = e_i^T M\tilde{b} = (M\tilde{b})_i = \tilde{x}_i$, which shows that \tilde{x}_i is attained. Second, let $x'' = DM(\tilde{b} - 2(\tilde{x}_i/\beta_i)\Delta e_i)$ and define a diagonal matrix D' by $D'_{ii} = -1$ and $D'_{jj} = D_{jj}$ otherwise. Then $(I - D\Delta D')DM = DM - D\Delta(I - 2e_i e_i^T)M = DM - D(M - I) + 2D\Delta e_i e_i^T M = D + 2D\Delta e_i e_i^T M$, hence $(I - D\Delta D')x'' = (D + 2D\Delta e_i e_i^T M)(\tilde{b} - 2(\tilde{x}_i/\beta_i)\Delta e_i) = D\tilde{b} + 2\tilde{x}_i D\Delta e_i(-1/\beta_i + 1 - (2/\beta_i)(m_i - 1)) = D\tilde{b} = b_c + D\delta$, which shows that x'' is a solution to the system $(I - D\Delta D')x'' = b_c + D\delta$ where $|(I - D\Delta D') - I| = \Delta$, $|(b_c + D\delta) - b_c| = \delta$ and $x''_i = e_i^T DM(\tilde{b} - 2(\tilde{x}_i/\beta_i)\Delta e_i) = \tilde{x}_i - 2(\tilde{x}_i/\beta_i)(m_i - 1) = \tilde{x}_i/\beta_i$. This proves that \tilde{x}_i/β_i is attained, hence also the upper bound $\max\{\tilde{x}_i, \tilde{x}_i/\beta_i\}$ in (9) is attained. The proof for the lower bound follows from the result just obtained by applying it to the case $b_c := -b_c$ as in the part 3). \square

The quantities r_i and h_i correct the influence of the approximate inverses R and M ; they vanish if $R = A_c^{-1}$ and $M = (I - G)^{-1} \geq 0$ are used. The last statement of the theorem is Hansen's optimality result [1] as reformulated in [2]. Matrices R and $M \geq 0$ satisfying (1) exist if and only if $\varrho(|A_c^{-1}|\Delta) < 1$ holds (Theorem 1 in [3]).

Acknowledgements

This work was supported by the Czech Republic Grant Agency under grant GA ČR 201/98/0222.

References

- 1 HANSEN, E. R.: Bounding the solution of interval linear equations. *SIAM J. Numer. Anal.* **29** (1992), 1493–1503.
- 2 ROHN, J.: Cheap and tight bounds: The recent result by E. Hansen can be made more efficient. *Interval Computations* **4** (1993), 13–21.
- 3 REX, G., ROHN, J.: A note on checking regularity of interval matrices. *Linear and Multilinear Algebra* **39** (1995), 259–262.

Addresses: JIŘÍ ROHN, Faculty of Mathematics and Physics, Charles University, Malostranské nám. 25, 118 00 Prague, and Institute of Computer Science, Academy of Sciences, Pod vodárenskou věží 2, 182 07 Prague, Czech Republic.