



národní  
úložiště  
šedé  
literatury

**A seven-years based characterization of aerosol light scattering properties at Central European rural site: Variability and source apportionment**

Suchánková, Lenka  
2022

Dostupný z <http://www.nusl.cz/ntk/nusl-508644>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 09.04.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní [nusl.cz](http://nusl.cz).

## A seven-years based characterization of aerosol light scattering properties at Central European rural site: Variability and source apportionment

*Student: Mgr. Lenka Suchánková*

*Supervisor: Prof. RNDr. Ivan Holoubek, CSc.*

*Supervising Expert: Ing. Vladimír Ždímal, Dr.*

Atmospheric aerosols have a significant impact on the radiative forcing of Earth's climate, directly through the aerosol-radiation interactions (ARIs) i.e., scattering or absorption of incoming solar and outgoing infrared radiation, or indirectly through the aerosol-cloud interactions (ACIs).<sup>1,2</sup> Although there are studies addressing aerosol impact on local climate,<sup>3-5</sup> the information on long-term measurements at rural background site is missing. Such studies are important for understanding of local sources and long-range transport of both anthropogenic and natural aerosols. Therefore, we focused on temporal variations of the total light scattering ( $\sigma_{sp}$ ) and backscattering ( $\sigma_{bsp}$ ) coefficients and associated optical properties such as the Ångström exponent (SAE), backscattering ratio ( $b$ ), and asymmetry factor ( $g$ ) at a rural background site National Atmospheric Observatory Košetice (NAOK; 49°34'20.787"N, 15°4'48.155"E) in Central Europe. We measured  $\sigma_{sp}$  and  $\sigma_{bsp}$  at 5 min resolution at three wavelengths (450, 550, and 700 nm) using the Integrating Nephelometer TSI 3563 (PM10 inlet). Measurements were performed from 16 August 2012 to 31 December 2019. The analysis has been performed using R software version 4.1.0. The preliminary results show that the overall trend for both  $\sigma_{sp}$  and  $\sigma_{bsp}$  is downward from 2012-2019; the slope of the median trend line was  $-2.50 \text{ Mm}^{-1}/\text{yr}$  and  $-0.18 \text{ Mm}^{-1}/\text{yr}$  at 550 nm, respectively.  $\text{SO}_2$  and  $\text{NO}_x$  concentrations were well correlated with  $\sigma_{sp}$  and  $\sigma_{bsp}$  throughout the period, confirming their contribution in the light-scattering enhancement.  $b$  had a positive slope of the median trend line ( $0.012/\text{yr}$ ), indicating more efficient cooling effect alongside lower aerosol loading. Both  $\sigma_{sp}$  and  $\sigma_{bsp}$  reached higher values in the cold seasons (median,  $46.8$  and  $5.9 \text{ Mm}^{-1}$ ) than in the summer (median,  $25.4$  and  $4.1 \text{ Mm}^{-1}$ ). This phenomenon is probably related to the higher aerosol loading in winter due to higher energy consumption, poorer dispersion of pollutants, and lower planetary boundary

layer. Elevated SAE observed in summer indicates smaller particles and corresponds to secondary organic aerosol. On the other hand, decreased SAE during winter corresponds to bigger particles (higher atmospheric stability and thus aerosol aging).

In addition, the sources of scattering aerosols, chemical composition, meteorological conditions, particle size distribution, and radiative forcing will be further investigated at NAOK to better understand the direct effects of aerosols on local climate.

This work was supported by the Ministry of Education, Youth and Sports of CR within the CzeCOS program, grant number LM2018123 within the National Sustainability Program I (NPU I), grant number LO1415, European Regional Development Fund-Project ACTRIS-CZ LM201818122, ACTRIS IMP ([www.actris.eu](http://www.actris.eu)) which is supported by the European Commission under the Horizon 2020 – Research and Innovation Framework Programme, H2020-INFRADEV-2019-2, Grant Agreement number: 871115.

#### References

1. IPCC, 2013. Summary for Policymakers. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change; Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., A. Nauels, Y. X., Bex, V., Midgley, P. M., Eds.; 2013
2. Ramanathan, V.; Crutzen, P. J.; Kiehl, J. T.; Rosenfeld, D. Atmosphere: Aerosols, Climate, and the Hydrological Cycle. *Science*. 2001, 294 (5549), 2119–2124.
3. Delene, D. J.; Ogren, J. A. Variability of Aerosol Optical Properties at Four North American Surface Monitoring Sites. *J. Atmos. Sci.* 2002, 59 (6), 1135–1150.
4. Luoma, K.; Virkkula, A.; Aalto, P.; Petäjä, T.; Kulmala, M. Over a 10-Year Record of Aerosol Optical Properties at SMEAR II. *Atmos. Chem. Phys.* 2019, 19 (17), 11363–11382.
5. Pandolfi, M.; Alados-Arboledas, L.; Alastuey, A. et al. A European Aerosol Phenomenology - 6: Scattering Properties of Atmospheric Aerosol Particles from 28 ACTRIS Sites. *Atmos. Chem. Phys.* 2018, 18 (11), 7877–7911.