



národní
úložiště
šedé
literatury

The Impact of the IMO-2020 Shipping Regulation on the Atmospheric Deposition of Sulfur around the Western English Channel and the Use of the V/Ni ratio as a Shipping Marker.

Windell, L.
2022

Dostupný z <http://www.nusl.cz/ntk/nusl-509009>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 27.09.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní nusl.cz .

THE IMPACT OF THE IMO-2020 SHIPPING REGULATION ON THE ATMOSPHERIC DEPOSITION OF SULFUR AROUND THE WESTERN ENGLISH CHANNEL AND THE USE OF THE V/Ni RATIO AS A SHIPPING MARKER

Laurence WINDELL^{1,3}, Thomas BELL^{1,2}, Caroline WHITE¹, Angela MILNE¹, Jaroslav SCHWARZ³, Simon USSHER¹

¹ Biogeochemistry Research Centre, University of Plymouth, Plymouth, UK, windell@icpf.cas.cz

² Plymouth Marine Laboratory, Plymouth, UK

³ Institute of Chemical Process Fundamentals of the CAS, Prague, Czech Republic

Keywords: IMO-2020 Global sulfur cap, Shipping emissions, Sulfur, V/Ni ratio

INTRODUCTION

Anthropogenic aerosols are shown to have a great impact on marine ecosystems and human health, with the shipping industry contributing as a significant source of these emissions, and annual growth on the rise. In the marine environment, shipping emissions often dominate atmospheric deposition of sulfur. With around 70% of shipping activity being carried out within 400 km of the coast, its contribution to atmospheric sulfur is of great interest. The International Maritime Organisation (IMO) introduced a regulation on 01/01/2020 restricting global marine fuel sulfur content from 3.5% to 0.5% w/w.

The vanadium/nickel (V/Ni) ratio has been widely used as a marker of shipping emissions, with the range of 2.5 to 4 indicating shipping activity. The regulation has called into question the viability of the earlier use of the V/Ni ratio.

EXPERIMENTAL SETUP

The present study investigated the changes in marine aerosol chemical character around Penlee Point Atmospheric Observatory (PPAO; 50°19' N, 4°11' W) to gain insight into the efficacy of the IMO-2020 regulation. The observatory in Southwest England is ideally located next to a shipping lane with high marine traffic. Aerosol filter samples were collected from 2020-2021 (n=53). Filter samples were water-leached and subsequent leachates analysed for major ions using Ion Chromatography (IC), and trace elements using Inductively Coupled Plasma - Mass Spectrometry (ICP-MS). Concentrations of V/Ni and calculated non-sea-salt sulfate (nss-SO₄²⁻) were compared to PPAO datasets from 2015-16 (n=56) and 2017-18 (n=51) respectively. Where possible, HYSPLIT air mass back trajectories were run (Stein et al., 2015).

RESULTS AND CONCLUSIONS

Trace element analysis showed a drop in median V from 2.91 pmol/m³ to 1.44 pmol/m³, and Ni increase from 0.95 pmol/m³ to 4.8 pmol/m³ with the decrease of V being in line with a similar post-IMO-2020 study in China (Guangyuan et al., 2021). Median V/Ni ratio post-IMO decreased from 3.3 to 0.28.

Post-IMO-2020 nss-SO₄²⁻ concentrations were significantly lower than pre-IMO-2020 values (n=109, p<0.05), which was observed throughout all classified air masses,

with total median nss-SO₄²⁻ concentration dropping from 1.35 µg/m³ to 0.33 µg/m³ (Fig 1.).




Figure 1. (a) Pre- and post-regulation nss-SO₄²⁻, (b) Pre- and post-regulation V/Ni ratios. Horizontal lines represent previous ratio range for shipping emissions.

Results from the major ion analysis suggest great differences in marine atmospheric nss-SO₄²⁻ post-regulation, pointing towards a noticeable decrease in nss-SO₄²⁻ around PPAO, potentially influenced by the S regulation. Trace element results indicate a drastic change in V/Ni ratios, far different from the generally agreed-upon range of 2.5 to 4. The results have presented a need to further investigate the viability of the V/Ni ratio as a marker of shipping emissions, where the previous ratio must be re-evaluated.

ACKNOWLEDGEMENT

The work was partially supported by MEYS of the Czech Republic under grant ACTRIS-CZ LM2018122.

REFERENCES

- Yu, G., Zhang, Y., Yang, F., He, B., Zhang, C., Zou, Z., ... & Chen, J., Dynamic Ni/V Ratio in the Ship-Emitted Particles Driven by Multiphase Fuel Oil Regulations in Coastal China. *Environ. Sci. Tech.*, 55(22), 15031-15039 (2021).
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J., Cohen, M. D., & Ngan, F., NOAA's HYSPLIT atmospheric transport and dispersion modeling system. *Bulletin of the American Meteorological Society*, 96(12), 2059-2077(2015).