

## Properties and use of high-temperature biochar

Sedmihradská, Anežka 2023 Dostupný z http://www.nusl.cz/ntk/nusl-526213

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL). Datum stažení: 30.07.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní nusl.cz .

## Properties and use of high-temperature biochar

Student: Ing. Anežka Sedmihradská Supervisor: doc. Ing. Michael Pohořelý, Ph.D.

Despite being a hot topic in environmental research, biochar and its properties have been mainly investigated from the perspectives of feedstock type and pyrolysis temperature. Few studies have explored the effect of particle size on biochar properties.<sup>1</sup> Even when the effect of particle size on crop efficiency is recognized, there is a lack of corresponding data and analysis.<sup>2–7</sup> Therefore, we thoroughly analyzed (proximate and ultimate analysis, SEM, XRD, Raman spectroscopy) four particle size fractions (< 0.5, 0.5-2, 2-5, > 5 mm) of high-temperature biochar from a gasification power plant, the sizes reflecting those of biochar actually used in agriculture. The < 0.5 mm fraction was significantly different in terms of carbon structure, and carbon and ash content; consequently, pH, electric conductivity and heavy metal content also differed greatly. The corresponding results for the other particle fractions were broadly similar to each other. Interestingly, for all fractions, the porosity and H/C ratio were independent of particle size, meaning that, regardless of size, the investigated biochar fractions are equally stable in soil. The 0.5–2 mm fraction was then selected to be used in different industrial or agricultural processes such as the anaerobic digestion of wastewater or livestock feeding. The effect of biochar on these processes is currently under investigation, providing only initial data so far.

## References

- 1. He, P.; Liu, Y.; Shao, L.; Zhang, H.; Lü, F. Particle Size Dependence of the Physicochemical Properties of Biochar. *Chemosphere* **2018**, *212*, 385–392.
- Prodana, M.; Silva, C.; Gravato, C.; Verheijen, F. G. A.; Keizer, J. J.; Soares, A. M. V. M.; Loureiro, S.; Bastos, A. C. Influence of Biochar Particle Size on Biota Responses. *Ecotoxicol. Environ. Saf.* 2019, 174, 120–128.
- de Jesus Duarte, S.; Glaser, B.; Pellegrino Cerri, C. Effect of Biochar Particle Size on Physical, Hydrological and Chemical Properties of Loamy and Sandy Tropical Soils. *Agronomy* 2019, 9 (4), 165.

32

- 4. Werdin, J.; Conn, R.; Fletcher, T. D.; Rayner, J. P.; Williams, N. S. G.; Farrell, C. Biochar Particle Size and Amendment Rate Are More Important for Water Retention and Weight of Green Roof Substrates than Differences in Feedstock Type. *Ecol .Eng.* **2021**, *171*, 106391.
- Alghamdi, A. G.; Alkhasha, A.; Ibrahim, H. M. Effect of Biochar Particle Size on Water Retention and Availability in a Sandy Loam Soil. *J. Saudi Chem. Soc*, 2020, 24 (12), 1042–1050.
- Głąb, T.; Palmowska, J.; Zaleski, T.; Gondek, K. Effect of Biochar Application on Soil Hydrological Properties and Physical Quality of Sandy Soil. *Geoderma* 2016, 281, 11–20.
- Hussain, R.; Ghosh, K. K.; Ravi, K. Influence of Biochar Particle Size on the Hydraulic Conductivity of Two Different Compacted Engineered Soils. *Biomass Convers. Biorefin.* 2021, 13 (2), 801–811.