



národní
úložiště
šedé
literatury

Liquid mixing time and gas holdup in a bubble column bioreactor

Terentyak, Mark
2023

Dostupný z <http://www.nusl.cz/ntk/nusl-526219>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 11.04.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní [nusl.cz](http://www.nusl.cz).

Liquid mixing time and gas holdup in a bubble column bioreactor


Student: Ing. Mark Terentyak

Supervisor: Ing. Mária Zedníková, Ph.D.

Supervising Expert: Sandra Cristina Kordač Petronilho Orvalho, Ph.D.

Bioreactors are widely used across many sectors of industry, especially in pharmacy and food production. Bubble columns are one of the bioreactor types suitable for cultivation of organisms sensitive to high shear forces, which often require careful supply of oxygen and other important nutrients. Therefore, controlled aeration and homogenization need to be established, especially in the case of tall bubble columns with high aspect ratios (liquid height to column width).¹ The aim of the research is the experimental study of homogenisation and aeration in a bubble column in dependence on operational conditions and aspect ratio.

Experiments were carried out in a bubble column with an inner diameter of 0.19 m, gas flow rate in the range 2–20 m³/h and aspect ratios ranging in 3.0–6.0. Aeration was quantified by gas holdup measured from the change of dispersion height. Homogenisation was quantified by mixing time t_{95} measured via the colorimetric method based on tracer experiments. Acid and alkaline solutions were used



as tracers and concentration changes were visualized by a pH indicator. Colour changes were recorded by camera and evaluated via image processing.²

Obtained results provide a significant influence of aspect ratio on both gas holdup and t_{95} . Based on data analysis, a correlation between t_{95} and gas holdup was developed considering an influence of operation conditions and aspect ratio.

The research is supported by the Internal grant agency of the Institute of Chemical Process Fundamentals of the CAS and by the Czech Academy of Sciences and the German Academic Exchange Service through the Mobility plus project DAAD-21-05.

References

1. Garcia-Ochoa, F.; Gomez, E. Bioreactor scale-up and oxygen transfer rate in microbial processes: an overview. *Biotechnol. Adv.* **2009**, 27 (2), 153–176.
2. Rosseburg, A.; Fitschen, J.; Wutz, J.; Wucherpfennig, T.; Schlüter, M. Hydrodynamic inhomogeneities in large scale stirred tanks – Influence on mixing time. *Chem. Eng. Sci.* **2018**, 188, 208–220.