



národní  
úložiště  
šedé  
literatury

## **Detection of Signed Homonuclear Couplings between Low Abundant Nuclei**

Blechta, Vratislav  
2011

Dostupný z <http://www.nusl.cz/ntk/nusl-71625>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 26.06.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní [nusl.cz](http://www.nusl.cz) .

## C-16

**DETECTION OF SIGNED HOMONUCLEAR COUPLINGS BETWEEN  
LOW ABUNDANT NUCLEI**

Vratislav Blechta and Jan Schraml

*Institute of Chemical Process Fundamentals,  
Academy of Sciences of the Czech Republic,  
Rozvojova 135, Prague 6 – Suchbát, 165 02*

Detection of signed homonuclear couplings between low abundant nuclei, e.g.  $J(\text{C},\text{C})$  or  $J(\text{Si},\text{Si})$ , is an uneasy task. Until now no measured signs of  ${}^nJ(\text{Si},\text{Si})$ ,  $n > 1$ , were published and signed  ${}^nJ(\text{C},\text{C})$ ,  $n > 1$ , are very rare.

Recently we published CSEc [1] method for detection of signs of  ${}^nJ(\text{Si},\text{C})$ ,  $n > 1$ , couplings applicable to other pairs of low abundant nuclei as well. The subject of this lecture is a modification of CSEc method for detection of signed homonuclear couplings. The new sequence is neither trivial nor simple application of the CSEc but some tricks are used to design broadband pulse sequence without selective pulses.

The sequence was proved on a few samples by measuring carbon-carbon two- and three-bond couplings. For the first time signs of  ${}^2J(\text{Si}-\text{O}-\text{Si})$  couplings were measured on a set of siloxane oligomers with different bond maps. The results will be presented.

Financial support provided by The Grant Agency of the AS CR (grant no. IAA 400720706) is gratefully acknowledged.