



národní
úložiště
šedé
literatury

A not-a-priori-exponential necessary and sufficient condition for regularity of interval matrices

Hooshyarbakhsh, V.
2012

Dostupný z <http://www.nusl.cz/ntk/nusl-80997>

Dílo je chráněno podle autorského zákona č. 121/2000 Sb.

Tento dokument byl stažen z Národního úložiště šedé literatury (NUŠL).

Datum stažení: 30.09.2024

Další dokumenty můžete najít prostřednictvím vyhledávacího rozhraní nusl.cz .



Institute of Computer Science
Academy of Sciences of the Czech Republic

**A not-a-priori-exponential necessary
and sufficient condition for
regularity of interval matrices**

Vahideh Hooshyarbakhsh, Raena Farhadsefat and
Jiri Rohn

Technical report No. V-1147

02.01.2012



Institute of Computer Science
Academy of Sciences of the Czech Republic

A not-a-priori-exponential necessary and sufficient condition for regularity of interval matrices


Vahideh Hooshyarbakhsh,¹ Raena Farhadsefat² and Jiri Rohn³

Technical report No. V-1147

02.01.2012

Abstract:

We describe a not-a-priori-exponential necessary and sufficient condition for regularity of interval matrices which is an easy consequence of an earlier result on interval linear equations.



Keywords:

Interval matrix, regularity, necessary and sufficient condition.⁴

¹Department of Applied Mathematics and Computer Sciences, Hamedan Branch, Islamic Azad University, Hamedan, Iran, e-mail: vhoshiar@iauh.ac.ir.

²Young Researchers Club, Hamedan Branch, Islamic Azad University, Hamedan, Iran, e-mail: rfarhad@iauh.ac.ir.

³Institute of Computer Science, Czech Academy of Sciences, Prague, Czech Republic, e-mail: rohn@cs.cas.cz. This author's work was supported by the Institutional Research Plan AV0Z10300504.

⁴Above: logo of interval computations and related areas (depiction of the solution set of the system $[2, 4]x_1 + [-2, 1]x_2 = [-2, 2]$, $[-1, 2]x_1 + [2, 4]x_2 = [-2, 2]$ (Barth and Nuding [1])).

Checking regularity of interval matrices is a known NP-hard problem. Forty necessary and sufficient regularity conditions are summed up in [2]; all of them are exponential because they explicitly or implicitly contain the quantifier “for each $z \in \{-1, 1\}^n$ ”. The condition given below is, to these authors’ knowledge, the *first* ever published not-a-priori-exponential regularity condition because instead of $\{-1, 1\}^n$ it employs only a subset Z of it. Cardinality of the set Z varies with the data, but its minimal value is 1. Notation used: e_j is the j th column of the $n \times n$ identity matrix, $e = (1, 1, \dots, 1)^T \in \mathbb{R}^n$, $\text{diag}(z)$ is the $n \times n$ diagonal matrix with diagonal vector z and for an $x \in \mathbb{R}^n$, $\text{sgn}(x)$ is defined by $(\text{sgn}(x))_i = 1$ if $x_i \geq 0$ and $(\text{sgn}(x))_i = -1$ otherwise.

Theorem 1. *An $n \times n$ interval matrix $\mathbf{A} = [A_c - \Delta, A_c + \Delta]$ is regular if and only if A_c is nonsingular and there exists a subset Z of $\{-1, 1\}^n$ having the following properties:*

(a) $\text{sgn}(A_c^{-1}e) \in Z$,

(b) for each $z \in Z$ the inequalities

$$(QA_c - I) \text{diag}(z) \geq |Q|\Delta, \quad (0.1)$$

$$(QA_c - I) \text{diag}(-z) \geq |Q|\Delta \quad (0.2)$$

have matrix solutions Q_z and Q_{-z} , respectively,

(c) if $z \in Z$, $Q_{-z}e \leq Q_z e$, and $(Q_{-z}e)_j(Q_z e)_j \leq 0$ for some j , then $z - 2z_j e_j \in Z$.

Proof. “If”: The assumptions (a)-(c) imply that the three assumptions of Theorem 3 in [3] are met for the system of interval linear equations $\mathbf{A}x = [e, e]$ whose solution set in virtue of the same theorem is bounded, hence \mathbf{A} is regular. “Only if”: If \mathbf{A} is regular, then (a) and (c) are satisfied for $Z = \{-1, 1\}^n$ and for each $z \in \{-1, 1\}^n$ the equations

$$(QA_c - I) \text{diag}(z) = |Q|\Delta,$$

$$(QA_c - I) \text{diag}(-z) = |Q|\Delta$$

have (even unique) solutions, see [2]. ▀

Hence we can also formulate the theorem in the following way:

Theorem 2. *An $n \times n$ interval matrix $\mathbf{A} = [A_c - \Delta, A_c + \Delta]$ is regular if and only if A_c is nonsingular and there exists a subset Z of $\{-1, 1\}^n$ having the following properties:*

(a) $\text{sgn}(A_c^{-1}e) \in Z$,

(b) for each $z \in Z$ the equations

$$(QA_c - I) \text{diag}(z) = |Q|\Delta, \quad (0.3)$$

$$(QA_c - I) \text{diag}(-z) = |Q|\Delta \quad (0.4)$$

have matrix solutions Q_z and Q_{-z} , respectively,

(c) if $z \in Z$, $Q_{-z}e \leq Q_z e$, and $(Q_{-z}e)_j(Q_z e)_j \leq 0$ for some j , then $z - 2z_j e_j \in Z$.

Notice that if $z \in \{-1, 1\}^n$, then $z - 2z_j e_j \in \{-1, 1\}^n$ (in (c)), so that $Z \subseteq \{-1, 1\}^n$; thus Z is defined recursively by (a) and (c). In practical computations, equations (0.3), (0.4) are solved instead of inequalities (0.1), (0.2) as it was done in the function `qzmatrix`, using the subfunction `absvaleqn`, in [4].

Bibliography

- [1] W. Barth and E. Nuding, *Optimale Lösung von Intervallgleichungssystemen*, Computing, 12 (1974), pp. 117–125. [1](#)
- [2] J. Rohn, *Forty necessary and sufficient conditions for regularity of interval matrices: A survey*, Electronic Journal of Linear Algebra, 18 (2009), pp. 500–512. http://www.math.technion.ac.il/iic/ela/ela-articles/articles/vol18_pp500-512.pdf. [2](#)
- [3] J. Rohn, *A general method for enclosing solutions of interval linear equations*, Technical Report 1067, Institute of Computer Science, Academy of Sciences of the Czech Republic, Prague, March 2010. <http://uivtx.cs.cas.cz/~rohn/publist/genmeth.pdf>. [2](#)
- [4] J. Rohn, *An algorithm for computing the hull of the solution set of interval linear equations*, Linear Algebra and Its Applications, 435 (2011), pp. 193–201. [2](#)